



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁴ : C12Q 1/70, 1/68, C12N 15/00 C12N 7/00, 1/20</p>	<p>A1</p>	<p>(11) International Publication Number: WO 89/ 02935 (43) International Publication Date: 6 April 1989 (06.04.89)</p>
<p>(21) International Application Number: PCT/US88/03399 (22) International Filing Date: 29 September 1988 (29.09.88) (31) Priority Application Numbers: 102,180 247,017 (32) Priority Dates: 29 September 1987 (29.09.87) 20 September 1988 (20.09.88) (33) Priority Country: US (71) Applicant: PRAXIS BIOLOGICS, INC. [US/US]; 30 Corporate Woods, Rochester, NY 14623-1493 (US). (72) Inventors: PARADISO, Peter, R. ; 6 Guilford Way, Pittsford, NY 14534 (US). HILDRETH, Stephen, W. ; 159 Willowbend Road, Rochester, NY 14618 (US). HU, Branda, T. ; 9 Hadley Court, Pittsford, NY 14534 (US). MARTIN-GALLARDO, Antonia ; 873 Pinna- cle Road, Pittsford, NY 14534 (US). ARUMUGH- AM, Rasappa ; 7 Cactus Drive, West Henrietta, NY 14586 (US).</p>	<p>(74) Agent: MISROCK, S., Leslie; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US). (81) Designated States: AT (European patent), AU, BE (Eu- ropean patent), CH (European patent), DE (Euro- pean patent), DK, FR (European patent), GB (Euro- pean patent), IT (European patent), JP, KR, LU (Eu- ropean patent), NL (European patent), SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: RESPIRATORY SYNCYTIAL VIRUS: VACCINES AND DIAGNOSTIC ASSAYS</p>		
<p>(57) Abstract</p> <p>Polypeptides, nucleotides, and compositions useful for preparing diagnostic reagents for and vaccines against hu- man Respiratory Syncytial Virus are disclosed. The polypeptides include short polypeptides which are related to a neutral- izing and fusion epitope of the Respiratory Syncytial Virus fusion protein or a neutralizing epitope of the G protein.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

RESPIRATORY SYNCYTIAL VIRUS: VACCINES AND DIAGNOSTIC ASSAYS

1. FIELD OF THE INVENTION

Respiratory Syncytial (RS) virus is a major cause of lower respiratory disease in infancy and early childhood.

5 It is a matter of great medical and scientific interest to provide safe and effective vaccines against this virus.

RS virus is an enveloped RNA virus. The major outer envelope proteins, the F protein (also known as the fusion protein or fusion glycoprotein) and the G protein, play a
10 key role in RS viral infection because antibodies directed against these proteins can neutralize the virus.

For the purposes of this Application, a neutralizing epitope on a viral protein is an epitope which is essential for virus infectivity as defined by the fact that antibody
15 binding to the epitope neutralizes the virus. Likewise, a fusion epitope on a viral protein is an epitope which is essential for virus-cell fusion or intercellular spread of infection by fusion between an infected-cell and an uninfected cell as defined by the fact that antibody-binding
20 to the epitope abrogates fusion.

The present invention relates to compositions and methods of preparation of proteins and polypeptides associated with the outer envelope of RS virus. More particularly, one embodiment of the invention is directed to
25 compositions and methods for preparation of proteins and polypeptides related to the fusion protein of RS virus. The proteins and polypeptides of this embodiment of the invention are related to a neutralizing epitope, or a fusion epitope, or both, of the fusion protein, and may be used as
30 immunogens in vaccine formulations including multivalent vaccines for active immunization and for the generation of antibodies for use in passive immunization, as well as reagents for diagnostic assays.

-2-

Another embodiment of the invention is directed to compositions and methods for preparation of proteins and polypeptides related to the G protein of RS virus. The proteins and polypeptides of this embodiment of the invention are related to a neutralizing epitope of the G protein, and may be used as immunogens in vaccine formulations including multivalent vaccines for active immunization and for the generation of antibodies for use in passive immunization, as well as reagents for diagnostic assays.

The novel proteins and polypeptides related to a neutralizing epitope, a fusion epitope, or both can be obtained by using either recombinant DNA or chemical synthetic methods. Additionally, the invention relates to novel DNA sequences and vectors useful for expressing RS virus related proteins and polypeptides and to cells which harbor the novel DNA sequences and vectors.

It should be noted that an epitope is a three dimensional structure generated by the molecular arrangement of an underlying molecular entity. In the present invention, the underlying molecular entities are polypeptides. It is well known that the structural properties of polypeptides of which three dimensional configuration is one, may only be minutely changed by the introduction of a small number of modifications such as substitutions, insertions and deletions of one or more amino acids. Generally, such substitutions in the amino acid sequence of a polypeptide are in the amount of less than twenty percent, more usually less than ten percent. Generally, conservative substitutions are less likely to make significant structural changes than non-conservative substitutions, which in turn are less likely to make significant structural changes than insertions or deletions.

35

-3-

Examples of conservative substitutions are glycine for alanine; valine for isoleucine; aspartic acid for glutamic acid; asparagine for glutamine; serine for threonine; lysine for arginine; phenylalanine for threonine; and the converse of the above. Therefore, it is to be understood that the present invention embraces modified polypeptides so long as the epitope is unchanged.

It is also well known that viral epitopes may exhibit strain-to-strain variations. Adjustment by the above-indicated modifications may indeed be used advantageously.

Finally, the fusion protein and G protein related polypeptides or proteins of the invention, like the bonafide viral proteins, may be labeled or unlabeled, bound to a surface, conjugated to a carrier, and the like, depending on the use to which they are put.

2. BACKGROUND OF THE INVENTION

2.1. RESPIRATORY SYNCYTIAL VIRUS IN DISEASE

RS virus is a major cause of lower respiratory disease in infancy and early childhood (McIntosh and Chanock, 1985, in Virology, Fields, B. (ed), Raven, NY, pp. 1285-1304). In all geographical areas, it is the major cause of bronchiolitis and pneumonia in infants and young children. The agent reinfects frequently during childhood, but illness produced by reinfection is generally milder than that associated with the initial infection and rarely causes major problems.

RS virus is an enveloped RNA virus of the family Paramyxoviridae and of the genus pneumovirus. The two major envelope proteins are the G protein, which is responsible for attachment of the virus to the host cell membrane, and the fusion protein, which is responsible for fusing the

-4-

virus and cell membranes. Virus-cell fusion is a necessary step for infection. Fusion protein is also required for cell-cell fusion which is another way to spread the infection from an infected cell to an uninfected cell.

5 Antibodies directed against the fusion protein or against the G protein can neutralize the virus. However, only antibodies to the fusion protein will block the spread of the virus between cells, i.e. have anti-fusion activity. Thus, antibodies to the fusion protein will protect against
10 circulating virus as well as inhibit the spread, between cells, of an established infection. Antibodies to the fusion protein (both polyclonal antisera against purified fusion protein and monoclonal antibodies which contain both neutralizing and anti-fusion activity) have been found to be
15 protective in animal models against infection (Walsh et al., 1984, Infect. Immun. 43:756-758).

2.2. IMMUNOLOGICAL APPROACH TO THE PREVENTION OF RS VIRUS INFECTION

20 A practical means for protection of infants and young children against upper and lower respiratory disease would be protective vaccination against RS virus. Vaccination of expectant mothers (active immunization) would protect young children by passive transfer of immunity, either trans-placentally, or through the mother's milk. Several
25 approaches to an RS virus vaccine are possible, but some of them have proven unsuccessful in the past.

Vaccination with killed RS virus vaccine has been tried and found to be ineffective (Kim et al., 1969, Am. J. Epid. 89:422). Not only were children not protected, but in
30 some cases, subsequent infections with RS virus resulted in atypical and more severe disease than in the unimmunized controls. This phenomenon is not unique to RS virus and has

-5-

been seen also in killed paramyxovirus vaccines such as measles. It has been suggested that the reason for the failure of the past inactivated RS virus vaccine was due to inactivation of the biologically functional epitopes on either or both of the viral envelope glycoproteins. That is to say, the neutralizing and fusion epitopes on the killed virus vaccine were "denatured". As a result, the vaccinated subject did not experience the biologically functional neutralizing and fusion epitopes. Therefore, when the vaccinated subject encountered a live virus, the resultant antibody response did not yield protective immunity. Instead, there was an antibody mediated inflammatory response which often resulted in a more severe disease (Choppin and Scheid, 1980, Rev. Inf. Dis., 2:40-61).

The second approach to an RS virus vaccine has been to attenuate live virus. Temperature sensitive mutants (Wright *et al.*, 1982, Infect. Immun. 37:397-400) and passage attenuated virus (Belshe *et al.*, 1982, J. Inf. Dis. 145:311-319) have proven to be poorly infectious and not efficacious in the prevention of disease when used as immunogens in RS virus vaccines. However, in these cases, there was no atypical disease as a result of vaccination.

Based on our current knowledge of the structure of RS virus and the immune response to infection, it is clear that a useful vaccine to this virus must be effective in inducing production of antibodies to the fusion protein and/or the G protein. Of particular importance to protective immunity is the production of antibodies that inhibit fusion and therefore, can stop the spread of virus between cells in the respiratory tract. Additionally, it may be helpful to induce a cell mediated immune response, including the stimulation of cytotoxic T cells (CTL's) which are useful against RS virus infected cells. The various vaccine

-6-

formulations of the present invention are directed to meeting both these objectives.

2.3. RECOMBINANT DNA TECHNOLOGY AND GENE EXPRESSION

5 Recombinant DNA technology involves insertion of specific DNA sequences into a DNA vehicle (vector) to form a recombinant DNA molecule which is capable of being replicated in a host cell. Generally, but not necessarily, the inserted DNA sequence is foreign to the recipient DNA
10 vehicle, i.e., the inserted DNA sequence and DNA vector are derived from organisms which do not exchange genetic information which in nature, or the inserted DNA sequence comprises information which may be wholly or partially artificial. Several general methods have been developed
15 which enable construction of recombinant DNA molecules. For example, U.S. Pat. No. 4,237,224 to Cohen and Boyer describes production of such recombinant plasmids using processes of cleavage of DNA with restriction enzymes and joining the DNA pieces by known methods of ligation.

20 These recombinant plasmids are then introduced by means of transformation and replicated in unicellular cultures including procaryotic organisms and eucaryotic cells grown in tissue culture. Because of the general applicability of the techniques described therein, U.S. Pat. No. 4,237,224 is hereby incorporated by reference into the
25 present specification. Another method for introducing recombinant DNA molecules into unicellular organisms is described by Collins and Hohn in U.S. Pat. No. 4,304,863 which is also incorporated herein by reference. This method
30 utilizes a packaging, transduction system with bacteriophage vectors (cosmids).

DNA sequences may also be inserted into viruses, for example, vaccinia virus. Such recombinant viruses may be

-7-

generated, for example, by transfection of plasmids into cells infected with virus (Chakrabarti et al., 1985, Mol. Cell. Biol. 5:3403-3409).

Regardless of the method used for construction, the
5 recombinant DNA molecule is preferably compatible with the host cell, i.e., capable of being replicated in the host cell either as part of the host chromosomes or as an extra-chromosomal element. The recombinant DNA molecule or recombinant virus preferably has a marker function which
10 allows the selection of the desired recombinant DNA molecule(s) or virus(es). In addition, if all of the proper replication, transcription and translation signals are correctly arranged on the recombinant DNA molecule, the foreign gene will be properly expressed in the transformed
15 or transfected host cells.

Different genetic signals and processing events control gene expression at different levels. For instance, DNA transcription is one level, and messenger RNA (mRNA) translation is another. Transcription of DNA is dependent
20 upon the presence of a promoter which is a DNA sequence that directs the binding of RNA polymerase and thereby promotes RNA synthesis. The DNA sequences of eucaryotic promoters differ from those of procaryotic promoters. Furthermore, eucaryotic promoters and accompanying genetic signals may
25 not be recognized in or may not function in a procaryotic system.

Similarly, translation of mRNA in procaryotes depends upon the presence of the proper procaryotic signals which differ from those of eucaryotes. Efficient translation of
30 mRNA in procaryotes requires a ribosome binding site called the Shine-Dalgarno (SD) sequence on the mRNA. For a review on maximizing gene expression, see Roberts and Lauer, 1979, Methods in Enzymology 68:473.

-8-

Many other factors complicate the expression of foreign genes in procaryotes even after the proper signals are inserted and appropriately positioned. One such factor is the presence of an active proteolytic system in E. coli and other bacteria. This protein-degrading system appears to destroy foreign proteins selectively. A tremendous utility, therefore, would be afforded by the development of a means to protect eucaryotic proteins expressed in bacteria from proteolytic degradation. One strategy is to construct hybrid genes in which the foreign sequence is ligated in phase (i.e., in the correct reading frame) with a procaryotic structural gene. Expression of this hybrid gene results in a recombinant protein product (a protein that is a hybrid of procaryotic and foreign amino acid sequences).

Similar considerations of gene expression in eukaryotic systems have been discussed in *Enhancers & Eukaryotic Gene Expression*, Gluzman & Shenk (Eds.), Cold Spring Harbor Laboratories, Cold Spring Harbor, New York 1983, and *Eukaryotic Viral Vectors*, Gluzman (Ed.), Cold Spring Harbor Laboratories, Cold Spring Harbor, New York 1982.

Successful expression of a cloned gene requires efficient transcription of DNA, translation of the mRNA and in some instances post-translational modification of the protein. Expression vectors have been developed to increase protein production from the cloned gene. In expression vectors, the cloned gene is often placed next to a strong promoter which is controllable so that transcription can be turned on when necessary. Cells can be grown to a high density and then the promoter can be induced to increase the number of transcripts. These, if efficiently translated will result in high yields of protein. This is an especially valuable system if the foreign protein is

-9-

deleterious to the host cell.

Several recombinant DNA expression systems are described below for the purpose of illustration only, and these examples should not be construed to limit the scope of the present invention.

2.3.1. E. COLI AS AN EXPRESSION VECTOR

Many E. coli plasmids are known and have been used to express foreign genes. For economic reasons, it would be highly preferable to be able to obtain a high level of expression. One way to obtain large amounts of a given gene product is to clone a gene on a plasmid which has a very high copy number within the bacterial cell. By increasing the number of copies of a particular gene, mRNA levels would normally also increase, which in turn would lead to increased production of the desired protein.

2.3.2. VACCINIA VIRUS AS AN EXPRESSION VECTOR

Vaccinia virus may be used as a cloning and expression vector. The virus contains a linear double-stranded DNA genome of approximately 187 kb pairs and replicates within the cytoplasm of infected cells. These viruses contain a complete transcriptional enzyme system (including capping, methylating and polyadenylating enzymes) within the virus core. This system is necessary for virus infectivity because vaccinia virus transcriptional regulatory sequences (promoters) allow for initiation of transcription by vaccinia RNA polymerase, but not by cellular RNA polymerase.

Expression of foreign DNA in recombinant viruses requires the fusion of vaccinia promoters to protein coding sequences of the foreign gene. Plasmid vectors, also called insertion vectors have been constructed to insert the chimeric gene into vaccinia virus. One type of insertion

-10-

vector comprises: (1) a vaccinia virus promoter including the transcriptional initiation site; (2) several unique restriction endonuclease cloning sites downstream from the transcriptional start site for insertion of foreign DNA fragments; (3) nonessential vaccinia virus DNA (such as the thymidine kinase gene) flanking the promoter and cloning sites which direct insertion of the chimeric gene into the homologous nonessential region of the virus genome; and (4) a bacterial origin of replication and antibiotic resistance marker for replication and selection in E. coli. Examples of such vectors are described by MacKett (1984, J. Virol. 49:857-864).

Recombinant viruses are produced by transfection of recombinant bacterial insertion vectors containing the foreign gene into cells infected with vaccinia virus. Homologous recombination takes place within the infected cells and results in the insertion of the foreign gene into the viral genome. Immunological techniques, DNA plaque hybridization, or genetic selection can be used to identify and isolate the desired recombinant virus. These vaccinia recombinants retain the functions essential for infectivity and can be constructed to accommodate up to approximately 35 kb of foreign DNA.

Expression of a foreign gene can be detected by enzymatic or immunological assays (e.g., immunoprecipitation, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay, or immunoblotting). Additionally, naturally occurring membrane glycoproteins produced from recombinant vaccinia infected cells are glycosylated and may be transported to the cell surface. High expression levels can be obtained by using strong promoters or cloning multiple copies of a single gene.

-11-

2.3.3. BACULOVIRUS AS AN EXPRESSION VECTOR

A baculovirus, such as Autographica californica nuclear polyhedris virus (AcNPV) has also been used as a cloning or expression vector. The infectious form of AcNPV
5 is normally found in a viral occlusion. This structure is largely composed of polyhedrin polypeptide in which virus particles are embedded. Polyhedrin gene expression occurs very late in the infection cycle, after mature virus particles are formed. Therefore, polyhedrin gene expression
10 is a dispensable function, i.e., non-occluded virus particles produced in the absence of polyhedrin gene expression are fully active and are capable of infecting cells in culture. According to European Patent Application Serial No. 84105841.5 by Smith et al., a recombinant
15 baculovirus expression vector can be prepared in two steps. First, baculovirus DNA is cleaved to produce a fragment comprising a polyhedrin gene or a portion thereof, which fragment is inserted into a cloning vehicle. The gene to be expressed is also inserted into the cloning vehicle; and it
20 is so inserted that it is under control of the polyhedrin gene promoter. This recombinant molecule is called a recombinant transfer vector. Normally, the recombinant transfer vector is amplified in appropriate host cells. Second, the recombinant transfer vector formed in this way
25 is mixed with baculovirus helper DNA and used to transfect insect cells in culture to effect recombination and incorporation of the cloned gene at the polyhedrin gene locus of the baculovirus genome. The resultant recombinant baculovirus is used to infect susceptible insects or
30 cultured insect cells.

3. SUMMARY OF THE INVENTION

The present invention is directed to polypeptides and proteins related to a neutralizing epitope, a fusion epitope, or both, of respiratory syncytial (RS) virus glycoproteins, including the fusion protein and G protein, as well as molecularly cloned gene or gene fragments encoding such polypeptides and proteins.

One embodiment of the present invention is directed to polypeptides and proteins related to a neutralizing epitope, a fusion epitope, or both, of the fusion protein of respiratory RS virus as well as molecularly cloned genes or gene fragments, which encode these polypeptides or proteins. Another embodiment of the present invention is directed to polypeptides and proteins related to a neutralizing epitope of the G protein of RS virus as well as molecularly cloned genes or gene fragments, which encode these polypeptides or proteins. The polypeptides or proteins of the present invention may be used as immunogens in subunit vaccine formulations for RS virus or as reagents in diagnostic immunoassays for RS virus. The polypeptides or proteins of the present invention may be produced using recombinant DNA techniques or synthesized by chemical methods.

The present invention is also directed to methods for the molecular cloning of and expression of genes or gene fragments encoding a neutralizing or a fusion epitope, or both, of RS virus fusion protein or a neutralizing epitope of the G protein. Accordingly, the invention is also directed to the construction of novel polynucleotide sequences and their insertion into vectors, including both RNA and DNA vectors, to form recombinant molecules or viruses which can be used to direct expression of polypeptides or proteins related to the epitope(s) in the appropriate host cells. The vectors include plasmid DNA, viral DNA, human viruses, animal viruses, insect viruses and bacterial phages.

-13-

Recombinant viruses or extracts of cells which comprise the polypeptides or proteins of the present invention can also be used as immunogens in viral vaccine formulations. Since a fusion or neutralizing epitope of a virus will be recognized as "foreign" in the host animal, humoral and cell mediated immune responses directed against the epitope(s) will be induced. In a properly prepared vaccine formulation, this should protect the host against subsequent RS virus infection.

10 The polypeptides and proteins of the present invention can also be used as reagents in immunoassays such as ELISA tests and radioimmunoassays to detect RS virus infections in blood samples, body fluids, tissues, and so forth.

15 The polynucleotide sequences of the present invention can also be used as reagents in nucleic acid hybridization assays to detect RS virus in blood samples, body fluids, tissues, and so forth.

4. BRIEF DESCRIPTION OF THE FIGURES

20 Figure 1 shows the complete nucleotide and amino acid sequences of the RS virus fusion protein, reproduced from Collins et al., 1985, Proc. Nat'l. Acad. Sci. USA, 81:7683-7687.

25 Figure 2 shows the electrophoretic behavior in SDS-polyacrylamide gel of 1 ug of the purified RS virus fusion protein under various conditions. In lane 2, the protein was reduced with 5% beta-mercaptoethanol and heated to 100°C prior to electrophoresis; in lane 3, the protein was heated to 100°C prior to electrophoresis; in lane 4, the protein was not heated or reduced prior to electrophoresis. Lane 1 contains standard marker proteins whose molecular weights are indicated in the left margin. The right margin shows the molecular weight of the various fusion protein components.

-14-

The gel was stained with silver for visualization.

Figure 3 shows the positions of synthetic polypeptides 1, 2, 3, 4, and 5 (spl - sp5) on a linear map of the F₁ subunit. The F₁ subunit encompasses amino acids 137-574.

5 Figure 3 also shows the exact amino acid sequence of the synthetic polypeptides.

Figure 4 shows the positions of proteolytic fragments of the fusion protein on a linear map of the F₁ subunit.

Figure 5 is a diagrammatic representation of an E. coli recombinant vector containing the complete nucleotide
10 sequence of the RS virus fusion protein gene.

Figure 6 shows the amino and carboxy terminal sequences of several recombinant proteins expressed in E. coli.

15 Figure 7 shows the positions of the recombinant proteins on a linear map of the F₁ subunit. The fragments shown in shaded bars were reactive with the L4 monoclonal antibody and those shown in solid bars were non-reactive with the L4 monoclonal antibody.

20 Figure 8 illustrates a dot blot autoradiogram of the four (4) synthetic polypeptides defined by the chains of amino acids shown in the left margin. Lanes 1-4 contain 20 ug, 15 ug, 10 ug and 5 ug of peptide respectively. Lane 5 contains a positive fusion protein control. The blot was
25 reacted with the L4 monoclonal antibody and then with ¹²⁵I-protein A.

Figure 9 is a diagrammatic representation of a recombinant expression vector of pPX1044 containing the complete nucleotide sequence of the RS virus G protein gene.

30

5. DETAILED DESCRIPTION OF THE INVENTION

According to one embodiment of the present invention, a region of the RS virus fusion protein which defines a

35

-15-

neutralizing and fusion epitope[s] has been identified. A method for producing novel proteins, polypeptides and peptides comprising the epitope[s], and the polynucleotide sequences which encode such novel protein and polypeptides
5 are provided.

The fusion protein of RS virus has an apparent molecular weight of 70,000 daltons by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). The primary (cistronic) translation product consisting of 574
10 amino acid residues is synthesized and subsequently processed by enzymatic cleavage at a trypsin sensitive site yielding two subunits designated F_1 (amino acids 137-574) and F_2 (amino acids 1-136).

The complete nucleotide sequence of the fusion protein gene from the A2 strain of RS virus (Collins et al., 1984, Proc. Nat'l Acad. Sci. USA, 81:7683-7687) which encodes the fusion protein of 574 amino acids is illustrated in Figure 1. The numbering system depicted in Figure 1 is used throughout this application. The F_1 (apparent molecular
20 weight 48,000 dalton) and F_2 (apparent molecular weight 23,000 dalton) subunits are linked through a disulfide bond to form a protein which is designated $F_{1,2}$ (apparent molecular weight about 70,000 daltons). When purified from RS virus or infected cells, the native fusion protein exists
25 predominantly as a dimer (apparent molecular weight 140,000 daltons). The dimeric form is the most immunogenic form of the RS virus fusion protein.

Figure 2 shows the electrophoretic mobility of various fusion protein components using SDS-PAGE. In lane 2, the
30 protein was reduced with 5% beta-mercaptoethanol and heated to 100°C prior to electrophoresis; in lane 3, the protein was heated to 100°C prior to electrophoresis; in lane 4, the protein was not heated or reduced prior to electrophoresis.

-16-

Lane 1 contains standard marker proteins whose molecular weights are indicated in the left margin. The right margin shows the molecular weight of the various fusion protein components. The gel was stained with silver for visualization (Morrissey, 1981, Anal. Biochem. 117:307-310).

Active immunization of cotton rats with purified RS virus fusion protein (the 140,000 dalton form) results in the production of antibodies which are effective in virus neutralization and in preventing fusion (see, for example, Section 6.2, infra). As demonstrated in Section 6.2, this immunization protected the lung and nasal tissues from infection by RS virus. Similar results have been obtained in baboons (see, Section 6.3, infra). Additionally, active immunization of animals with purified RS virus fusion protein previously reduced with beta-mercaptoethanol protects animals from subsequent RS virus infection (see Section 6.2, infra).

According to another embodiment of the present invention, substantially pure polypeptides and proteins related to a neutralizing epitope of the RS virus G protein are provided. The RS virus G protein has an apparent molecular weight of about 84,000-90,000 daltons and is highly glycosylated. The nucleotide sequence of the gene encoding the RS virus G protein has been disclosed (Satake et al., 1985, Nucleic Acid Res. 13:7795-812; Wertz et al., 1985, Proc. Nat'l Acad. Sci. USA 82:4075-79). The entire gene sequence encoding the RS G protein has been cloned and expressed in a recombinant expression vector (see Section 9, infra). Surprisingly, active immunization of animals with a recombinant non-glycosylated RS virus G protein induced a protective immune response (see Section 10, infra).

A monoclonal antibody to the fusion protein,

-17-

designated L4 (Walsh and Hruska, 1983, J. Virol. 47:171-177), is capable of both neutralizing RS virus and inhibiting fusion. Therefore, it appears that this antibody reacts with an epitope on the fusion protein which is
5 essential to both infectivity and the fusion function of the fusion protein, i.e., this antibody reacts with both a fusion epitope and a neutralizing epitope. Moreover, passive transfer of the L4 monoclonal antibody will protect cotton rats from virus infection in their lungs (Walsh et
10 al., 1984, Infect. Immun., 43:756-758).

In addition to the L4 type antifusion, neutralizing epitope, there is yet another antifusion, neutralizing epitope of the fusion protein which may be conformationally dependent. This epitope is recognized by monoclonal antibody A5 (Walsh et
15 al., 1986, J. Gen. Virol. 67:505-513). The reactivity of this antibody appears to be dependent upon the native conformation of the fusion protein. When the conformation of the fusion protein is altered either by heat treatment or by heat treatment and the reduction of disulfide bonds, the A5 type
20 epitope is destroyed. The present studies have examined the role of L4 type and A5 type epitopes of the RS virus fusion protein on the formulation of the protein as an effective vaccine. Results from a competitive ELISA indicate that one third of the fusion protein specific antibody in human adults recovering from natural RS infection, is directed against
25 epitopes recognized by either L4 or A5. Vaccination with purified F protein produces a response which can be 80% blocked by a combination of L4 and A5. Competitive binding to the F protein, shows that these epitopes overlap by 40%.
30 Based on these studies, it appears that the functional antifusion, neutralizing epitopes can be classified into two categories, L4 type and A5 type.

5.1. IDENTIFICATION OF NEUTRALIZING AND/OR FUSION EPITOPE[S] OF RS VIRUS FUSION PROTEIN

According to the present invention, the region of the RS virus fusion protein which is an epitope responsible for eliciting both neutralizing and antifusion antibodies has now been determined. This region has been defined by three methods. The first method employs defined proteolytic cleavage of the native protein. The second method relates to cloning and expressing fragments of the fusion protein gene, for example, in *E. coli*. The third relates to synthesis of synthetic polypeptides which bind to neutralizing and antifusion antibodies. In all three methods reactivity with the L4 monoclonal antibody was used to identify desired fragments. Any other monoclonal antibody which is capable of neutralizing and preventing fusion of RS virus may be used.

5.1.1. MAPPING BY DEFINED PROTEOLYTIC CLEAVAGE

The L4 monoclonal antibody was tested for its ability to bind to the fusion protein subunits (F_1 and F_2) by protein immunoblot (Western blot) analysis. Using Western blot as described in Section 8.1, *infra*, the L4 monoclonal antibody was able to bind only to the F_1 subunit. Additionally, the L4 monoclonal antibody has been shown to bind to serotype B virus (Anderson et al., 1985, J. Inf. Dis. 151:623-33) defined by the prototype virus designated strain 18537.

In order to map the F_1 subunit, synthetic polypeptides are prepared which correspond to various regions along the F_1 subunit. These synthetic polypeptides are coupled to a carrier protein, such as keyhole lympet hemocyanin (KLH), and then used separately to immunize rabbits (see Section 8, *infra*). In the particular example exemplified in Section

-19-

8.2., five antisera (anti-sp1 through anti-sp5) were obtained. Each antiserum produced in the rabbits reacted with the uncoupled synthetic polypeptide corresponding to the immunogen which induced the antiserum, i.e., anti-sp1
5 reacted with sp1; anti-sp2 reacted with sp2, etc. All five antisera reacted with the F₁ subunit of the fusion protein.

The purified fusion protein is then subjected to proteolytic cleavage under a variety of conditions (see Section 8, infra). In the example illustrated in Section
10 8.2, infra, the proteases used were (1) trypsin which specifically cleaves after lysine and arginine residues, (2) endoproteinase Arg-C which cleaves specifically after arginine residues, and (3) endoproteinase Lys-C which
15 cleaves specifically after lysine residues. Cleavage after a residue means the breaking of a peptide bond at the carboxyl end of the residue. It should be understood that other combinations of proteases can be used. Therefore, the presently exemplified combination should not be construed as a limitation on the present invention.

20 Cleavages are also performed in the presence and absence of the L4 monoclonal antibody. The cleaved protein fragments are separated by SDS-PAGE and the cleavage products analyzed by Western blot analysis for the ability to bind to the L4 monoclonal antibody as well as the anti-
25 synthetic polypeptide antibodies (See an exemplary illustration Section 8.2, infra). The positions of the proteolytic fragments within the fusion protein sequence are deduced from the reactivities of these cleavage fragments with each of the anti-synthetic polypeptide antisera. For
30 example, as illustrated in Section 8.2., a cleavage fragment generated by trypsin digestion which fragment reacted with anti-sp1 would be expected to comprise amino acids 155-169. Furthermore, the fragment must have arginine or lysine at

-20-

its carboxyl end and a N-terminal residue which follows a lysine or arginine in the sequence of Figure 1. Finally, the molecular weight of a fragment can be determined by its mobility in SDS-PAGE. This set of information uniquely
5 predicts that the 28,000 dalton fragment generated by trypsin digestion and reactive with anti-spl spans amino acids 137-394 of the F₁ polypeptide. The positions of the other cleavage fragments are similarly deduced.

The relationship between the positions of the cleavage
10 fragments and the reactivities of these fragments to the L₄ monoclonal antibody are analyzed. From the example presented in Section 8.2., it can readily be seen that amino acids 283-327 define the region which is common to all of the L4 positive fragments and absent from all of the L4
15 negative fragments. This region, therefore, comprises a virus neutralizing and/or antifusion epitope of the RS virus defined by the L4 monoclonal antibody.

5.1.2. CLONING AND EXPRESSING FRAGMENTS OF THE FUSION PROTEIN

20 The cDNA illustrated in Figure 1 containing the complete nucleotide sequence of the fusion protein gene is cloned into a cloning vector such as the E. coli plasmid vector pBR322. Due to the degeneracy of the nucleotide coding sequences, other DNA sequences which encode
25 substantially the same amino acid sequence as depicted in Figure 1 may be used. These include but are not limited to nucleotide sequences comprising all or portions of the fusion protein nucleotide sequences depicted in Figure 1
30 which are altered by the substitution of different codons that encode the same or a functionally equivalent amino acid residue within the sequence (for example, an amino acid of the same polarity) thus producing a silent change.

-21-

Regions of the fusion protein gene are excised from the cloning vector by restriction endonuclease digestion and ligated into a compatible expression vector (see infra). In the experimental example described in Section 8.2., infra,
5 the E. coli expression vector pUC19 (Yanish-Perron et al., 1985, Gene 33:103-19) was used. The expressed recombinant proteins are screened for reactivity first with polyclonal rabbit antiserum to native fusion protein to identify recombinant fragments and then with L4 monoclonal to
10 identify those fragments comprising the neutralizing and antifusion epitope. As illustrated in Section 8.3., the RS virus fusion protein sequence common to all of the recombinant proteins reactive with the L4 monoclonal antibody is defined by amino acid residues 253-298.

15

5.1.3. SYNTHESIS OF ANTIGENIC PEPTIDES

In order to confirm the identify of the neutralizing and antifusion epitope identified as described above, synthetic polypeptides can be prepared corresponding in
20 particular to amino acid residues 299-315; 294-315; 289-315; and 283-315 of the RS virus fusion protein. These peptides are analyzed for reactivity with L4 monoclonal antibody. As illustrated in Section 8.4., infra, polypeptides containing residues 294-315; 289-315; and 283-315 react positively with
25 L4; whereas peptide 299-315 does not. This indicates that a neutralizing and fusion epitope resides between residues 283-298 and may be as small as residues 294-299.

5.2. PREPARATION OF PROTEINS, POLYPEPTIDES AND PEPTIDES RELATED TO RS VIRUS FUSION PROTEIN AND G PROTEIN

30 The proteins, polypeptides and peptides of the present invention can be prepared in a wide variety of ways. The polypeptides, because of their relatively short size may be

35

-22-

synthetized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, 1984, Solid Phase Peptide Synthesis, 2d Ed., Pierce Chemical Co. The structural properties of polypeptides, of which three dimensional configuration is one, may only be minutely changed by the introduction of a small number of modifications such as substitutions, insertions and deletions of one or more amino acids. Generally, such substitutions in the amino acid sequence of a polypeptide are in the amount of less than twenty percent, more usually less than ten percent. Generally, conservative substitutions are less likely to make significant structural changes than non-conservative substitutions, which in turn are less likely to make significant structural changes than insertions or deletions. Examples of conservative substitutions are glycine for alanine; valine for isoleucine; aspartic acid for glutamic acid; asparagine for glutamine; serine for threonine; lysine for arginine; phenylalanine for threonine; and the converse of the above. Therefore, it is to be understood that the present invention embraces modified polypeptides so long as the epitope of the RS virus fusion protein remains unchanged.

It is also well known that viral epitopes may exhibit strain-to-strain variations. Adjustment by the above-indicated modifications may indeed be used advantageously.

The polypeptides of the present invention may be employed as labeled or unlabeled compounds depending on their use. By label is intended a moiety which provides, directly or indirectly, a detectable signal. Various labels may be employed, such as radionuclides, enzymes, fluorescers, chemilumescers, enzyme substrates, cofactors

-23-

or inhibitors, particles (e.g. magnetic particles), ligands (e.g. biotin) and receptors (e.g. avidin) or the like. In addition, the polypeptides may be modified in a variety of ways for binding to a surface e.g. microtiter plate, glass beads, chromatographic surface, e.g. paper, cellulose and the like. The particular manner in which the polypeptides are joined to another compound or surface is conventional and finds ample illustration in the literature. See, for example, U.S. Pat. Nos. 4,371,515; 4,487,715 and the patents cited therein.

Alternatively, recombinant DNA technology may be employed to prepare the polypeptides biosynthetically.

5.3. INSERTION OF THE RS VIRUS FUSION PROTEIN OR G PROTEIN CODING SEQUENCES INTO EXPRESSION VECTORS

The nucleotide sequence coding for the RS virus fusion protein or a portion thereof or for the RS virus G protein or a portion thereof is inserted into an appropriate expression vector, i.e. a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. According to a preferred embodiment of this mode of the invention, nucleotide sequences coding for a neutralizing and/or fusion epitope of the fusion protein is inserted into an appropriate expression vector. The coding sequence may be extended at either the 5' and 3' terminus or both termini to extend biosynthetically the polypeptide while retaining the epitope. The extension may provide an arm for linking, e.g., to a label (see below), to a carrier or surface. The extension may provide for immunogenicity which may otherwise be lacking in some of the shorter antigenic polypeptides of the invention.

A variety of host-vector systems may be utilized to

-24-

express the protein-coding sequence. These include but are not limited to mammalian cell cultures such as Chinese hamster ovary cell host cultures, etc.; mammalian cell systems infected with virus (e.g. vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g. baculovirus); microorganisms such as yeast containing yeast vectors or bacteria transformed with bacteriophage DNA, plasmid DNA or cosmid DNA. In one embodiment the expression vector can be an attenuated enteroinvasive bacteria including but not limited to Salmonella spp., enteroinvasive E. coli (EIEC), and Shigella spp. Such bacterium can invade gut epithelial tissue, disseminate throughout the reticuloendothelial system and gain access to mesenteric lymphoid tissue where they multiply and induce humoral and cell-mediated immunity. The expression elements of these vectors vary in their strength and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used. For instance, when cloning in mammalian cell systems, promoters isolated from the genome of mammalian cells, (e.g., mouse metallothionien promoter) or from viruses that grow in these cells, (e.g. vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted sequences.

Specific initiation signals are also required for efficient translation of inserted protein coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where the RS virus fusion protein gene or the RS virus G protein gene including its own initiation codon and adjacent sequences are inserted into the appropriate expression vectors, no additional translational control signals may be needed. However, in cases where only

-25-

a portion of the RS virus fusion protein or G protein coding sequence is inserted, exogenous translational control signals, including the ATG initiation codon must be provided. The initiation codon must furthermore be in phase
5 with the reading frame of the protein coding sequences to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be a variety of origins, both natural and synthetic.

Any of the methods previously described for the
10 insertion of DNA fragments into a vector may be used to construct expression vectors containing a chimeric gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and synthetic
15 techniques and in vivo recombinations (genetic recombination).

The invention is not limited to the use of E. coli or procaryotic expression vectors. Expression vectors which can be used include, but are not limited to the following
20 vectors or their derivatives: human or animal viruses such as vaccinia viruses or adenoviruses; insect viruses such as baculoviruses; yeast vectors; bacteriophage vectors, and plasmid and cosmid DNA vectors to name but a few.

In cases where an adenovirus is used as an expression
25 vector, the RS virus fusion protein gene or fragment thereof or the RS virus G protein gene or fragment thereof is ligated to an adenovirus transcriptional/translational control complex, e.g., the late promoter and tripartite leader sequences. This chimeric gene is then inserted in
30 the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the RS virus fusion

-26-

protein or G protein related protein in infected hosts. Presently, there are two strains of adenovirus (types 4 and 7) approved and used as vaccines for military personnel. They are prime candidates for use as vectors to express the RS virus fusion protein or G protein gene and fragments thereof.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the chimeric gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers, (e.g., zinc and cadmium ions for metallothionein promoters). Therefore, expression of the genetically engineered RS virus fusion protein or G protein or fragment thereof may be controlled. This is important if the protein product of the cloned foreign gene is lethal to host cells. Furthermore, modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products are important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

25

5.4. IDENTIFICATION OF RECOMBINANT EXPRESSION VECTORS CAPABLE OF REPLICATING AND EXPRESSING THE INSERTED GENE

Expression vectors containing foreign gene inserts can be identified by three general approaches: (a) DNA-DNA hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the first approach, the presence of a foreign gene inserted in

35

-27-

an expression vector can be detected by DNA-DNA hybridization using probes comprising sequences that are homologous to the foreign inserted gene. In the second approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus etc.) caused by the insertion of foreign genes in the vector. For example, if the RSV virus fusion protein gene or fragment thereof is inserted within the marker gene sequence of the vector, recombinants containing the RS virus fusion protein inserted can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the foreign gene product expressed by the recombinant. Such assays can be based on the physical, immunological, or functional properties of the gene product.

Once a particular recombinant DNA molecule is identified and isolated, several methods may be used to propagate it, depending on whether such a recombinant constitutes a self-replicating unit (a replicon). A self replicating unit, e.g., plasmids, viruses, cells etc., can multiply itself in the appropriate cellular environment and growth conditions. Recombinants lacking a self-replicating unit will have to be integrated into a molecule having such a unit in order to be propagated. For example, certain plasmid expression vectors upon introduction into a host cell need to be integrated into the cellular chromosome to ensure propagation and stable expression of the recombinant gene. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity.

-28-

5.5. IDENTIFICATION AND PURIFICATION OF THE EXPRESSED GENE PRODUCT

Once a recombinant which expresses the RS virus fusion protein gene or fragment thereof or the RS virus G protein or fragment thereof is identified, the gene product should be analyzed. This can be achieved by assays based on the physical, immunological or functional properties of the product. Immunological analysis is especially important where the ultimate goal is to use the gene products or recombinant viruses that express such products in vaccine formulations and/or as antigens in diagnostic immunoassays.

A variety of antisera are available for analyzing immunoreactivity of the product, including but not limited to L4 monoclonal antibody, A5 monoclonal antibody, polyclonal antisera raised against purified fusion protein or G protein as described in Section 6 or 9, infra.

The protein should be immunoreactive whether it results from the expression of the entire gene sequence, a portion of the gene sequence or from two or more gene sequences which are ligated to direct the production of chimeric proteins. This reactivity may be demonstrated by standard immunological techniques, such as radioimmunoprecipitation, radioimmune competition, or immunoblots.

Once the RS virus fusion or G protein related protein is identified, it may be isolated and purified by standard methods including chromatography (e.g., ion exchange, affinity, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

-29-

5.6. DETERMINATION OF THE IMMUNOPOTENCY OF THE RECOMBINANT PRODUCT

Immunopotency of the RS virus fusion or G protein related product can be determined by monitoring the immune response of test animals following immunization with the purified protein, synthetic peptide or protein. In cases where the RS virus fusion protein or G protein related protein is expressed by an infectious recombinant virus, the recombinant virus itself can be used to immunize test animals. Test animals may include but are not limited to mice, rats, rabbits, primates, and eventually human subjects. Methods of introduction of the immunogen may include oral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal or any other standard routes of immunizations. The immune response of the test subjects can be analysed by three approaches: (a) the reactivity of the resultant immune serum to authentic RS viral antigens, as assayed by known techniques, e.g., enzyme linked immunosorbant assay (ELISA), immunoblots, radioimmunoprecipitations, etc., (b) the ability of the immune serum to neutralize RS virus infectivity in vitro (see Section 6, infra), (c) the ability of the immune serum to inhibit virus fusion in vitro (see Section 6, infra) and (d) protection from RS virus infection (see Section 6, infra).

5.7. FORMULATION OF A VACCINE

Many methods may be used to administer the vaccine formulations described herein to an animal or a human. These include, but are not limited to: oral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous and intranasal routes. The secretory IgA antibodies produced by the mucosal associated lymphoid tissue may play a major role in protection against RS virus infection by preventing the initial interaction of the pathogens with

-30-

the mucosal surface, or by neutralizing the important epitopes of the pathogens that are involved in infection/or spreading of the disease. Stimulation of mucosal immune responses, including production of secretory IgA antibodies may be of
5 major importance in conferring protection against lower and upper respiratory tract infection. When a live recombinant virus vaccine formulation is used, it may be administered
via the natural route of infection of the parent wild-type virus which was used to make the recombinant virus in the
10 vaccine formulation.

5.7.1. SUBUNIT VACCINE FORMULATIONS

The proteins and polypeptides of the present invention related to a neutralizing and/or fusion epitope of the
15 fusion protein of RS virus are useful as immunogens in a subunit vaccine to protect against lower respiratory disease and other disease symptoms of RS virus infection. Subunit vaccines comprise solely the relevant immunogenic material necessary to immunize a host. Vaccines prepared from
20 genetically engineered immunogens, chemically synthesized immunogens and/or immunogens comprising authentic substantially pure RS virus fusion protein or fragments thereof alone or in combination with similarly prepared RS virus G protein or fragments thereof, which are capable of
25 eliciting a protective immune response are particularly advantageous because there is no risk of infection of the recipients.

Thus, the RS virus fusion protein and/or G protein related proteins and polypeptides can be purified from
30 recombinants that express the neutralizing and/or fusion epitopes. Such recombinants include any of the previously described bacterial transformants, yeast transformants, cultured cells infected with recombinant viruses or cultured

-31-

mammalian cells such as Chinese hamster ovary cells that express the RS virus fusion protein epitopes (see Section 5.3, supra). Additionally the recombinants include recombinant attenuated enterovasive bacteria containing a

5 DNA sequence which encodes a neutralizing and/or fusion epitope of Respiratory Syncytial Virus fusion protein or a neutralizing epitope of Respiratory Syncytial Virus G protein. Such recombinants are prepared using methods similar to those described in United States Patent

10 Application Serial No. 104,735. These recombinant attenuated enteroinvasive bacteria are particularly suited for oral vaccine formulations. The recombinant protein or polypeptides can comprise multiple copies of the epitope of interest.

15 Alternatively, the RS virus fusion protein and/or G protein related protein or polypeptide can be chemically synthesized (see Section 5.2, supra). In yet another alternative embodiment, the RS virus fusion protein related protein or polypeptide or G related protein can be isolated

20 in substantially pure form from RS virus or cultures of cells infected with RS virus (see, for example, Section 6.1, or Section 10, infra).

Regardless of the method of production, the RS virus fusion protein or G protein, related protein or polypeptide

25 is adjusted to an appropriate concentration and can be formulated with any suitable vaccine adjuvant. The polypeptides and proteins may generally be formulated at concentrations in the range of 0.1 ug to 100 ug per kg/host. Physiologically acceptable media may be used as carriers.

30 These include, but are not limited to: sterile water, saline, phosphate buffered saline and the like. Suitable adjuvants include, but are not limited to: surface active substances, e.g., hexadecylamine, octadecylamine, octadecyl

-32-

amino acid esters, lysolecithin, dimethyl-
dioctadecylammonium bromide, N,N-dioctadecyl-N'-N-bis(2-
hydroxyethyl-propane diamine), methoxyhexadecylglycerol, and
pluronic polyols; polyamines, e.g., pyran, dextran sulfate,
5 poly IC, polyacrylic acid, carbopol; peptides, e.g., muramyl
dipeptide, dimethylglycine, tuftsin; oil emulsions; and
mineral gels, e.g., aluminum hydroxide, aluminum phosphate,
etc. The immunogen may also be incorporated into liposomes
or conjugated to polysaccharides and/or other polymers for
10 use in a vaccine formulation.

In yet another embodiment of this mode of the
invention, the RS virus fusion protein related protein or
polypeptide is a hapten, i.e., a molecule which is antigenic
in that it reacts specifically or selectively with cognate
15 antibodies, but is not immunogenic in that it cannot elicit
an immune response. In such case, the hapten may be
covalently bound to a carrier or immunogenic molecule; for
example, a large protein such as protein serum albumin will
confer immunogenicity to the hapten coupled to it. The
20 hapten-carrier may be formulated for use as a subunit
vaccine.

The polypeptides and proteins of the present invention
may be used when linked to a soluble macromolecular carrier.
Preferably, the carrier and the polypeptides and proteins of
25 the present invention are in excess of five thousand daltons
after linking. More preferably, the carrier is in excess of
five kilodaltons. Preferably, the carrier is a polyamino
acid, either natural or synthetic, which is immunogenic in
animals, including humans. The manner of linking is
30 conventional. Many linking techniques are disclosed in U.S.
Pat. No. 4,629,783 which is incorporated herein by
reference. Many cross-linking agents are disclosed in
1986-87 Handbook And General Catalog, Pierce Chemical

35

-33-

Company, (Rockford, Illinois) pages 311 to 340, which pages are incorporated herein by reference.

In yet another embodiment of this mode of the invention the immunogen of the vaccine formulation comprises
5 a mixture of polypeptides and proteins related to the RS virus fusion protein and the G protein of one or more virus subtypes.

5.7.2. VIRAL VACCINE FORMULATIONS

10 Another purpose of the present invention is to provide either a live recombinant viral vaccine or an inactivated recombinant viral vaccine which is used to protect against lower respiratory infections and other disease symptoms of RS virus. To this end, recombinant viruses are prepared
15 that express RS virus fusion protein related epitopes (see Section 5.2, supra). Where the recombinant virus is infectious to the host to be immunized but does not cause disease, a live vaccine is preferred because multiplication in the host leads to a prolonged stimulus, therefore,
20 conferring substantially long-lasting immunity. The infectious recombinant virus when introduced into a host can express the RS virus fusion protein related protein or polypeptide fragment from its chimeric gene and thereby elicit an immune response against RS virus antigens. In
25 cases where such an immune response is protective against subsequent RS virus infection, the live recombinant virus, itself, may be used in a preventative vaccine against RS virus infection. Production of such recombinant virus may involve both in vitro (e.g., tissue culture cells) and in
30 vivo (e.g., natural host animal) systems. For instance, conventional methods for preparation and formulation of smallpox vaccine may be adapted for the formulation of live recombinant virus vaccine expressing an RS virus fusion protein related protein or polypeptide.

-34-

Multivalent live virus vaccines can be prepared from a single or a few infectious recombinant viruses that express epitopes of organisms that cause disease in addition to the epitopes of RS virus fusion protein. For example, a
5 vaccinia virus can be engineered to contain coding sequences for other epitopes in addition to those of RS virus fusion protein. Such a recombinant virus itself can be used as the immunogen in a multivalent vaccine. Alternatively, a
10 mixture of vaccinia or other viruses, each expressing a different gene encoding for an epitope of RS virus fusion protein and an epitope of another disease causing organism can be formulated in a multivalent vaccine.

Whether or not the recombinant virus is infectious to the host to be immunized, an inactivated virus vaccine
15 formulation may be prepared. Inactivated vaccines are "dead" in the sense that their infectivity has been destroyed, usually by chemical treatment (e.g., formaldehyde). Ideally, the infectivity of the virus is destroyed without affecting the proteins which are related
20 to immunogenicity of the virus. In order to prepare inactivated vaccines, large quantities of the recombinant virus expressing the RS virus fusion protein related protein or polypeptide must be grown in culture to provide the necessary quantity of relevant antigens. A mixture of
25 inactivated viruses which express different epitopes may be used for the formulation of "multivalent" vaccines. In certain instances, these "multivalent" inactivated vaccines may be preferable to live vaccine formulation because of potential difficulties with mutual interference of live
30 viruses administered together. In either case, the inactivated recombinant virus or mixture of viruses should be formulated in a suitable adjuvant in order to enhance the immunological response to the antigens. Suitable adjuvants

-35-

include, but are not limited to: surface active substances, e.g., hexadecylamine, octadecyl amino acid esters, octadecylamine, lysolecithin, dimethyl-dioctadecylammonium bromide, N, N-dioctadecyl-N'-N-bis (2-hydroxyethyl-propane
5 diamine), methoxyhexadecylglycerol, and pluronic polyols; polyamines, e.g., pyran, dextransulfate, poly IC, polyacrylic acid, carbopol; peptides, e.g., muramyl dipeptide, dimethylglycine, tuftsin; oil emulsions; and mineral gels, e.g., aluminum hydroxide, aluminum phosphate,
10 etc.

5.7.3. PASSIVE IMMUNITY AND ANTI-IDIOTYPIC ANTIBODIES

Instead of actively immunizing with viral or subunit vaccines, it is possible to confer short-term protection to
15 a host by the administration of pre-formed antibody against an epitope of RS virus fusion protein or an epitope of the RS virus G protein. Thus, the vaccine formulations can be used to produce antibodies for use in passive immunotherapy. Human immunoglobulin is preferred in human medicine because
20 a heterologous immunoglobulin may provoke an immune response to its foreign immunogenic components. Such passive immunization could be used on an emergency basis for immediate protection of unimmunized individuals exposed to special risks, e.g., young children exposed to contact with
25 RS virus patients. Alternatively, these antibodies can be used in the production of anti-idiotypic antibody, which in turn can be used as an antigen to stimulate an immune response against RS virus fusion protein or G protein epitopes.

30

5.8. DIAGNOSTIC ASSAYS

Yet another purpose of the present invention is to provide reagents for use in diagnostic assays for the

35

-36-

detection of antigens of the RS virus fusion protein or G protein (and hence RS virus) or for the detection of antibodies to the RS virus fusion protein or G protein in various body fluids of individuals suspected of RS virus infection.

5.8.1. IMMUNOASSAYS

In one mode of this embodiment, the RS virus fusion protein or G protein related proteins, polypeptides and peptides of the present invention may be used as antigens in immunoassays for the detection of RS virus in various patient tissues and body fluids including, but not limited to: blood, spinal fluid, sputum, nasal secretions, secretions of the respiratory tract, etc.

The proteins, polypeptides and peptides of the present invention may be used in any immunoassay system known in the art including, but not limited to: radioimmunoassays, ELISA assay, "sandwich" assays, precipitin reactions, gel diffusion immunodiffusion assays, agglutination assays, fluorescent immunoassays, protein A immunoassays and immunoelectrophoresis assays, to name but a few. U.S. Patent No. 4,629,783 and patents cited therein also describe suitable assays.

5.8.2. NUCLEIC ACID HYBRIDIZATION ASSAY

In another mode of this embodiment, the novel nucleotide sequence of the gene or gene fragment encoding the RS virus fusion protein related proteins polypeptides and peptides of the present invention may be used as probes in nucleic acid hybridization assays for the detection of RS virus in various patient body fluids, including but not limited to: blood, sputum, nasal secretions, secretions of the respiratory tract, etc.

35

-37-

The nucleotide sequences of the present invention may be used in any nucleic acid hybridization assay system known in the art including, but not limited to: Southern blots (Southern, 1975, J. Mol. Biol. 98:508); Northern blots (Thomas et al., 1980, Proc. Nat'l Acad. Sci. USA 77:5201-05); colony blots (Grunstein et al., 1975, Proc. Nat'l Acad. Sci. USA 72:3961-65), etc.

The following Examples are offered for the purpose of illustrating the present invention and are not to be construed to limit the scope of the present invention.

6. PROTECTION OF ANIMALS: RS VIRUS FUSION PROTEIN

6.1. GENERAL PROCEDURES

6.1.1. ISOLATION OF FUSION PROTEIN

Substantially pure RS virus fusion protein suitable for use as an immunogen in a subunit vaccine formulation was prepared essentially according to the procedure of Walsh et al. (1985, J. Gen. Virol. 66: 409-15). Purified protein has been prepared from three different virus strains including Long, A2 and 18537 and from two different cell lines, HEP-2 (ATCC No. CCL23) and Vero (ATCC No. CCL81). The protein derived from all sources is highly immunogenic and when used as an immunogen produces antibody which is virus neutralizing and antifusing. The electrophoretic behavior of the purified protein derived from the Long strain of virus in HEP-2 cells can be seen in Figure 2 under a variety of denaturing conditions. The protein is substantially pure of contaminating viral or cellular proteins.

The fusion glycoprotein purified as described above was found to have lipid covalently associated with the F₁ subunit. This was demonstrated by infecting Vero cells with

-38-

the A2 strain of RS virus and labeling the cultures with [9,10-³H]-palmitic acid. The purified protein was demonstrated to have ³H-palmitic acid associated with the 140,000 dalton dimeric form of the protein based on PAGE and
5 autoradiography. Furthermore, treatment of the fusion protein with heat alone or with heat plus reduction of disulfide bonds showed that the ³H-palmitic acid was associated with the 70,000 and 48,000 dalton (F₁ subunit) forms of the protein.

10 When ³H-palmitic acid labeled protein was extracted with chloroform:methanol (2:1 v/v), nearly all of the label remained associated with the protein and was not extracted as free lipid into the chloroform:methanol. Therefore, the palmitic acid is covalently attached. When the protein was
15 first treated with 1 M hydroxylamine at pH 7.0, the palmitic acid was then extracted into chloroform:methanol. Thus the protein-lipid bond is broken by hydroxylamine showing a covalent ester linkage. Since the bond was broken at neutral pH, a thioester linkage through cysteine on the
20 protein is the most likely bond although other ester bonds could be formed.

In addition, preliminary experiments suggest that myristic acid is also present in the RS virus F Protein. Purified F protein was hydrolysed in 1M methanolic HCl at
25 80°C for 24 hours; released lipid was extracted in hexane and analyzed by gas chromatography (GC) (Perkin Elmer 8500) using a fused silica column bonded with methyl silicone. The GC spectra obtained indicated the presence of myristic acid on the purified F protein.

30 The carbohydrate nature of the purified fusion protein obtained as described above was also characterized. The purified protein derived from the Long strain of RS virus obtained from infected HEP-2 cells was analyzed as follows:

-39-

The purified protein was methanolized with HCl-containing methanol, fully acetylated, O-deacetylated, and finally per-O-(trimethylsilyl)ated. The sugar residues were identified and quantitated by gas liquid chromatography (Reinhold, 1972, Methods in Enzymology, 25:244-49). The protein was found to have 5.75% total carbohydrate by weight. The sugar composition by percentage of total sugar was: 11.5% fucose, 3.3% xylose, 26.2% mannose, 9.8% galactose, 9.8% glucose, and 39.3% N-acetyl-glucosamine.

10

6.1.2. ASSAYS

6.1.2.1. VIRUS NEUTRALIZATION ASSAY

Virus neutralization assays were performed as follows:

15 Test serum samples and the positive control serum were heat inactivated at 56°C for 30 min. Test samples were serially diluted. All sera were then diluted with an equal volume containing about 50 plaque forming units (PFU) of RS virus, and incubated at 37°C for one hour. A pool of adult
20 sera which had previously been characterized by enzyme immunoassay, neutralization and antifusion assays was used for positive control. Sera which had previously been characterized and was known to be non-immune was used as negative control.

25 Each incubated serum-virus mixture was inoculated to HEp-2 cells (ATCC No. CCL23) in a separate well of 24 well plates and virus adsorption was allowed to take place for 2 hours at 37°C. The inocula were removed. The cell monolayers were washed and overlaid with modified Eagle's
30 medium plus 5% fetal bovine serum and 1% Sephadex®, and incubated at 37°C for 3 days. The overlay medium was removed and the cells were washed with phosphate buffered saline (PBS).

35

-40-

One ml of chilled PBS-methanol (1:5) solution was added to each well, and the cells were fixed for 30 min. at room temperature. The PBS-methanol fixative was removed, and one ml per well of 5% Carnation[®] instant milk in PBS, pH 6.8 (BLOTTO) was added. The plate was incubated for 30 minutes at 37°C.

The BLOTTO was removed. One-half ml per well of monoclonal antibodies against RS virus (previously titered and diluted with BLOTTO to a working concentration) was added, and the plate was incubated at 37°C for 1 hour. The antibodies were removed, and the fixed cells were washed twice with BLOTTO, 30 minutes each time.

One-half ml/well of horseradish peroxidase conjugated goat anti-mouse IgG (diluted 1:250 in BLOTTO) was added and the plate was incubated for 1 hour at 37°C. The goat antibodies were removed, and the fixed cells were again washed twice with BLOTTO, 30 minutes each time.

One-half ml/well of a peroxidase substrate solution (0.05% 4 chloro-1-naphthol, 0.09% H₂O₂ in PBS pH, 6.8) was added, and color was allowed to develop for 15-30 minutes at room temperature. The substrate solution was removed, and the wells were washed with water and air dried. The number of plaques in each well was determined.

The neutralization ability of a test serum sample is expressed as the dilution which results in a 60% reduction in plaque formation when compared to non-immune control serum expressed per ml of serum.

6.1.2.2. ANTI-FUSION ASSAY

Anti-fusion assays were performed as follows:

HEp-2 cells in 48 well plates were infected with RS virus at 25 PFU/well for 6 hours at 37°C. After infection, the culture medium was replaced with fresh culture medium

-41-

containing 0.1% K6 monoclonal antibodies (sterile filtered ascites fluid) and either a heat-inactivated test serum sample or a heat-inactivated non-immune control serum. K6 monoclonal antibody is specific for G glycoprotein of RS virus. The presence of K6 in the culture medium prevents virus spread via the culture medium. Sephadex® was added to a final concentration of 1% and the plate incubated for 3 days at 37°C. The culture medium was removed and the cells were washed with PBS.

One ml of chilled PBS-methanol (1:5) solution was added to each well, and the cells were fixed for 30 min. at room temperature. The fixative was removed, and one ml per well of BLOTTO was added. The plate was incubated for 30 minutes at 37°C.

The BLOTTO was removed. One-half ml/well of monoclonal antibodies against RS virus (previously titered and diluted with BLOTTO to a working concentration) was added, and the plate was incubated for 1 hr. at 37°C. The antibody solution was removed and the fixed cells were washed twice with BLOTTO, 30 minutes each time.

One-half ml per well of horseradish peroxidase conjugated goat anti-mouse IgG diluted 1:250 in BLOTTO was added, and the plate was incubated for 1 hour at 37°C. The goat antibodies were removed, and the fixed cells were again washed twice with BLOTTO, 30 minutes each time.

One-half ml per well of a peroxidase substrate solution (PBS pH 6.8 containing 0.05% 4 chloro-1-naphthol, 0.09% of H_2O_2) was added, and color was allowed to develop for 15-30 minutes at room temperature. The substrate solution was removed, and the wells were washed with water and air dried.

The number and typical size of plaques in the well corresponding to the non-immune control serum sample were

-42-

determined. The number of plaques of a similar size was then determined for the wells corresponding to the test serum samples. The anti-fusion titer of a test serum sample is expressed as the dilution which yields a 60% reduction in plaques scored when compared to non-immune control serum expressed per ml of serum.

6.1.2.3. ENZYME IMMUNOASSAY (EIA)

Antibody titer in serum samples was determined using an Enzyme Immunoassay (EIA) performed as follows:

RS virus fusion protein was diluted to 200 ng/ml in carbonate-bicarbonate buffer, pH 9.6. One hundred ul of the diluted antigen was added to each well of rows B-G of a flat-bottomed, 96 well Nunc[™] assay plate. In rows A and H, 100 ul of carbonate-bicarbonate buffer alone was added to each well. The plate was covered and incubated for 2 hours at 37°C with shaking and then stored overnight at 4°C to immobilize the antigen.

The supernatants were removed from the Nunc[™] assay plate and the plate was washed with 0.1% Tween/PBS pH 7.4 and pat dried.

Three antibody samples were assayed on each plate. Each sample was first diluted to a primary dilution in 0.2% Tween, 0.01 M EDTA/PBS pH 7.5 (0.2% TWN). The primary dilutions were further serially diluted as follows in a 96 well U-bottomed Falcon[™] plate:

(a) The primary dilutions of the samples were inoculated into row 2 at 200 ul/well. Sample 1 was inoculated in triplicate, e.g., in wells A2, B2, and C2; Sample 2 in duplicate e.g., in wells D2, E2; Sample 3 in triplicate e.g. in wells F2, G2, and H2.

(b) 100 ul of 0.2% TWN were inoculated into each well of rows 3-12.

-43-

(c) Serial dilutions were created by transferring sequentially 100 ul from a well in row 2 to the corresponding well in row 3 (e.g. B2 to B3; C2 to C3), a well in row 3 to the corresponding well in row 4, until row 12 was reached.

(d) To row 1, 100 ul of 0.2% TWN was added to each well as control.

One hundred ul of the primary dilutions were transferred from each well of the Falcon[™] plate to the corresponding well in the Nunc[™] plate, e.g., A2 (Falcon[™]) to A2 (Nunc[™]). The Nunc[™] assay plate was covered and incubated for 1 hour at 37°C with shaking. The supernatants were removed from the assay plate, and the plate was washed with 0.1% Tween/PBS and pat dried.

Goat anti-Mouse IgG alkaline phosphatase conjugate (TAGO[™]) was diluted with 0.3% Tween/PBS pH 7.0 (0.3% TWN) to a working dilution, e.g., 1:1500. The diluted conjugate (100 ul) was added to each well in rows 2-12. To row 1, 100 ul of 0.3% TWN were added to each well as control. The plate was covered and incubated for 1 hour at 37°C with shaking. The inocula was then removed, and the plate was washed with 0.1% Tween/PBS pH 7.4 and pat dried.

To each and every well, 100 ul substrate solution, 1 mg/ml in diethanolamine buffer pH 9.8 (SIGMA-104[™]) were added. The enzymatic reaction was allowed to take place at room temperature for 1 hour. The reaction was then stopped by adding 100 ul of 3N NaOH to each well. The extent of enzymatic reaction was determined by reading the optical density at 410 nm.

Rows A and H served as negative controls because no antigen was present; row 1 was also served as a negative control because no antibodies were present.

-44-

6.2. PROTECTION OF ANIMALS: HOMOLOGOUS AND HETEROLOGOUS PROTECTION

In one experiment, eighteen cotton rats were divided into 6 groups of 3 animals each. At week 0, 2 and 4-5, animals were actively immunized by intramuscular injection of 10 ug RS virus fusion protein (140,000 daltons), except Group 6 which received only 5 ug protein, in different adjuvants: Group 1, PBS; Group 2, IFA; Group 3, ISCOM; and Group 4, alum. Group 5 received intramuscular injections of PBS alone and served as the control group. Group 6 received intramuscular injections of RS virus fusion protein previously reduced using beta-mercaptoethanol, but without any adjuvant.

Serology assays were performed as described in Section 6.1.2, supra on serum samples obtained at week 6-7. Results are presented in Table 1.

Animals were challenged between weeks 6-7 with 1×10^4 PFU RS virus. Lungs were harvested on day 4 post-challenge. The presence and/or quantity of RS virus in lung tissues was determined. At day 4 post-infection, the lung and nasal turbinates were removed from the animals and homogenized in 1-2 ml of virus transport media (MEM, 5% FBS, 2 mM glutamine, 20 mM HEPES, 10% SPG). After centrifugation, the supernatants were serially diluted and applied to HEP-2 cells in 24 well tissue culture plates and the virus grown. Plaques were identified as described in Section 6.1.2.1. (Virus Neutralization Assay) and expressed per gram of tissue. Results are also presented in Table 1.

30

35

-45-

Table 1
IMMUNOGENICITY OF RSV FUSION
PROTEIN IN COTTON RATS

5	Group No. (Adjuvant) ^a	Serology ^b			Protection ^c	
		EIA	Virus Neutral- ization	Anti- fusion	Virus Present	Titer
10	Group 1 (PBS)	6.2	4.4	2.0	0	0
	Group 2 (IFA)	7.3	5.1	2.9	0	0
	Group 3 (ISCOM)	7.4	4.5	2.6	0	0
15	Group 4 (ALUM)	7.1	5.2	2.7	0	0
	Group 5 (control)	<1.0	<2.0	<1.0	3	4.8
20	Group 6 (none)	4.9	2.3	2.4	3	1.1

^a Group 6 received fusion protein treated with beta mercaptoethanol and no adjuvant.

^b Assays were performed on samples from week 7. Results represent the geometric mean titer three animals expressed in log₁₀ units. See text for experimental details.

^c Three animals in each of Groups 1-5 and 4 animals in Group 6 were challenged at weeks 6-7 and lungs were harvested on day 4 post-challenge. Virus was isolated from the lung tissues. "Virus present" represents the number of animals in each group in which virus was detectable. "Titer" represents the geometric mean titer (PFU/gm of tissue) of virus isolated from lung tissue.

-46-

As demonstrated in Table 1, immunization with RS virus fusion protein elicited production of antibodies effective virus neutralization and in preventing fusion. Moreover, results illustrated in Table 1 clearly demonstrate that lung tissues of immunized animals are effectively protected against subsequent RS virus infection.

When the fusion protein is treated (reduced) with beta-mercaptoethanol, the subunits are dissociated yielding free F_1 and F_2 . When cotton rats were immunized with 5 ug of the reduced protein without adjuvant, they also produced antibody that was neutralizing and had antifusion activity (see Table 1, Group 6), although the level of these activities was reduced. Similarly, the lungs of these animals were substantially protected from infection, however, with less efficacy. Thus, the subunits of the fusion glycoprotein, presented in a dissociated form to the immune system, are able to produce protective immunity. It should also be noted that under these conditions, the F_1 subunit, which contains the hydrophobic membrane anchor region, will aggregate forming higher molecular weight multimers which may enhance immunity.

Human RS virus is subdivided into two subtypes, A and B. The fusion proteins from both subtypes share antigenic determinants and these determinants are highly conserved among RS virus strains. Hence a series of experiments were conducted to determine whether immunization with F protein from one subtype of human RS virus would confer protection against infection by both subtypes of human RS virus.

In this series of experiments, 30 cotton rats were divided into 4 groups of experimental animals and 4 groups of control animals. At week 0, 2 and 4, experimental animals were actively immunized by intramuscular injection of 10 ug human RS virus fusion protein (140,000 daltons) obtained from RS virus A2 strain, a subtype A virus. Control animals were

-47-

similarly immunized with a placebo immunogen, i.e. PBS.

Serology assays were performed as described in Section 6.1.2, supra, to test the virus neutralization and anti-fusion capabilities of the induced antibodies against both A and B subtypes of human RS virus. Immunization with subtype A RS virus fusion protein induced antibodies which exhibited similar neutralization and anti-fusion activities against both subtypes A and B of RS virus (data not shown).

All experimental and control animals were challenged intranasally at week 6 with human RS virus as follows: Group 1 received strain A2 (subtype A) 6.2 log 10 PFU; Group 2, Long strain (subtype A) 6.1 log 10 PFU; Group 3, strain 9320 (subtype B) 3.5 log 10 PFU; and Group 4, strain 18537 (subtype B) 5.0 log 10 PFU. At 4 days post-challenge, the lungs were harvested and the virus titer was determined as described above. Results are presented in Table 2.

Table 2

PROTECTIVE EFFICACY OF FUSION PROTEIN
VACCINE AGAINST VARIOUS SUBTYPES OF RSV

Group No. ^a	Challenge Virus (Subtype)	Lung Titer (GMT) ^b	
		Experimental ^c	Control ^d
1	A2 (A)	1.1	5.0
2	Long (A)	1.1	4.0*
3	9320 (B)	1.43	5.1*
4	18537 (B)	1.1	3.5

^a Experimental animals in each group were immunized at week 0, 2 and 4 with 10 ug of F protein purified from RS virus strain A2, subtype A. Control animals in each group similarly received PBS as immunogen. All animals were challenged at week 6 with RS virus. See text for experimental details.

-48-

- b "Lung Titer (GMT)" represents the geometric mean titer (PFU/gm of tissue) of virus isolated from lung tissue.
- c N = 5.
- d N = 3, except * which designated N = 2.
- 5

As demonstrated in Table 2, immunization with fusion protein from a subtype A virus strain, induced significant protection against both a homologous subtype A virus and a heterologous sybtype B virus in all cases. Thus it is clear that immunization with F protein from one subtype RS virus confers protection against both subtypes of human RS virus.

10

6.3. PROTECTION OF BABOONS

Twelve juvenile baboons were divided into 3 groups of 4 animals each. Animals were injected intramuscularly with 20 ug purified RS virus fusion protein at one month intervals for 3 months as follows: Group 1 received immunogen in PBS; Group 2, immunogen in alum; and Group 3, PBS alone (control group).

15

20

Serology assays were performed two weeks after the third immunization as described in Section 6.1.2, supra. Results are presented in Table 3.

At 2 weeks post-immunization, 3-4 animals/group were challenged with 5.32 PFU of Strain A2 RS virus by direct inoculation into the lungs. Lung lavages were assayed at day 2 post-infection. Lung lavages were obtained by direct transfer of media into the lungs and collection by tube aspiration. The virus was titered as described in Section 6.2, supra. Results are also presented in Table 3.

25

30

-49-

Table 3

IMMUNOGENICITY OF RSV FUSION PROTEIN IN BABOONS

5	Group No. Immunogen(Adjuvant)	Serology ^a			Virus Assay ^b		
		EIA	Virus Neutral- ization	Anti fusion	+	-	TITER
10	Group 1 (PBS)	5.01	2.5	1.83	3	1	1.05
	Group 2 (ALUM)	6.0	3.34	2.56	0	4	0.00
	Group 3 (control)	<1.0	<2.0	<1.0	3	0	2.14

15 ^a Assays were performed two weeks after the last (third) immunization. Results represent the geometric mean titer 4 animals expressed in log₁₀ units. (See text for experimental details).

20 ^b Strain A2 RS virus (5.32 PFU) was inoculated directly into the lungs of 3-4 immunized animals in each group at two weeks post immunization. Lung lavages were assayed at day 2 post-infection. "+" represents the number of animals with detectable RS virus in lung lavages. "-" represents the number of animals without detectable RS virus in lung lavages. "Titer" is expressed as the geometric mean titer (PFU/gm) of tissue of virus isolated from lung tissue.

25 As shown in Table 3, immunization of juvenile baboons with purified RS virus elicited antibodies which were able to neutralize virus and prevent fusion. As further clearly demonstrated, immunized animals were protected against subsequent RS virus infection when alum was used as an
30 adjuvant (Group 2).

In another series of experiments, juvenile baboons were divided into a number of groups and immunized by intramuscular

-50-

injection of purified RS virus fusion protein as follows: 4 animals received 5 ug of immunogen in alum and 11 animals received 20 ug of immunogen in alum on days 0, 28 and 56; 6 animals received 100 ug of immunogen in alum on day 0, 20 ug on day 28 and 20 ug on day 56. Twelve animals received PBS alone on day 0, 28 and 56 and served as controls. All animals were challenged with $>10^6$ PFU of strain A2 RS virus on day 70 by direct inoculation into the lungs. Lung lavages were collected as described above on days 2, 3 and 4 post-challenge. Results from this series of experiments are summarized in Table 4.

Table 4

SUMMARY OF BABOON PROTECTION

Immunogen	N	RSV Isolation From Lung Lavages ^a		Titer
		(+)	(-)	
F (alum)	21	2	19	<0.7
PBS	12	12	0	3.0

^a "(+)" represents the number of baboons in which RS virus was detected in lung lavage. "(-)" represents the number of baboons in which no RS virus was detected in lung lavage. "Titer" is expressed as the geometric mean titer (PFU/ml of tissue) of virus isolated from lung lavage.

As summarized in Table 4, baboons were efficiently protected against RS virus infection by immunization with RS virus fusion protein. During the course of immunization and subsequent challenge experiments, blood chemistries and hematologic assays showed no sign of adverse reactions throughout the course of the experiments.

-51-

6.4. PROTECTION AGAINST BOVINE RS VIRUS

A crude preparation of human RS virus fusion protein (herein termed "crude human RS virus F protein" was prepared as follows: Vero cells infected with RS virus A2 strain were
5 extracted with a solution containing 50 mM Tris; 0.15 M NaCl; 1% Triton X-100 and 2% deoxycholate; pH 7.4 (Lysis Buffer). The F protein was obtained from the spent cell culture medium by solubilization of PEG pelleted virus using the Lysis Buffer. The resulting crude preparation was clarified by
10 centrifugation. An immunoaffinity purified preparation was prepared as described in Section 6.1.1.

Fifteen cows were divided into 5 groups of 3 animals each. Animals were injected intramuscularly with human RS virus fusion protein on days 0 and 21 as follows: Group 1
15 received crude human RS virus F protein (5 ug); Group 2, crude human RS virus F protein (20 ug); Group 3, purified human RS virus F protein, (20 ug); Group 4, a USDA approved bovine RS virus vaccine commercially available from Diamond Scientific Co. (Des Moines, Iowa) (2 ml on days 0 and 21) and Group 5,
20 PBS alone (control).

EIA's were performed on serum samples obtained on days 0, 10, 21 and 33 post-immunization as described in Section 6.1.2., supra, F protein from RS virus strain A2 as antigen. Results are presented in Table 5.

25

30

35

-52-

Table 5
IMMUNOGENICITY OF RS VIRUS FUSION PROTEIN IN COWS

Immunogen ^a	Serology (EIA) ^b			
	Day			
	0	10	21	33
5 HRSV F Protein (5 ug)*	-	-	3.6 (2/3)	5.8
HRSV F Protein (20 ug)*	-	3.8 (2/3)	4.2 (3/3)	6.2
HRSV F protein (20 ug)	-	3.8 (2/3)	4.6 (2/3)	5.9
Diamond vaccine (BRSV)	2.2 (1/3)	3.0 (1/3)	3.5 (2/3)	4.7
10 PBS	-	-	-	-

^a Cows were immunized at days 0 and 21 with either crude (*) or immunoaffinity purified F protein and the immunogenicity was compared with that of a commercially available USDA approved BRSV vaccine (Diamond Scientific Co. vaccine). See text for details.

^b Results represent the geometric mean titer of 3 animals expressed in log₁₀ units. The ratio in the parenthesis is the number of ELISA positive animals. "-" indicates that the titer was not detectable, i.e., less than 1.6.

20 The data presented in Table 5 shows that the human RS virus fusion protein obtained as described in Section 6.1 and after immunoaffinity purification elicits high levels of antibodies. Comparison with a commercially available bovine RS virus vaccine showed that the human RS virus F protein was
25 as immunogenic or more immunogenic in cows than the BRSV vaccine.

30 The protective immunogenicity of the human RS virus fusion protein against bovine RS virus was investigated in another series of experiments. Twenty five cotton rats were divided into 2 groups of 10 animals each and one group of 5 animals each. Animals were immunized intramuscularly on days 0 and 21 as follows: Group 1 received the commercially

35

-53-

available BRSV vaccine (Diamond); Group 2, human RS virus Long strain F and G protein (10 ug each)/Alum (affinity purified F and G proteins were obtained as described in Sections 6.1.1 and 10.1, respectively) ; and Group 3, PBS/Alum (control).

- 5 Serum samples were obtained and serological assays were performed as described in Section 6.2, supra, except that BRSV (strain 3758) was used for the neutralization and anti-fusion assays. Results are presented in Table 6.

10

Table 6

PROTECTIVE EFFICACY OF HUMAN RS
FUSION PROTEIN AGAINST BOVINE RS VIRUS^a

Immunogen	EIA		Serology ^b		Anti-fusion	
	Day		Neutralization		Day	
	21	28	21	28	21	28
Diamond	4.2 (10)	5.3 (6)	n.d.	2.6 (6)	n.d.	<1:20 (6)
F&G/Alum ^c	6.0 (10)	7.0 (8)	n.d.	4.3 (8)	n.d.	2.2 (8)
PBS/Alum	<1:100 (5)	<1:100 (10)	n.d.	<1:20 (4)	n.d.	<1:20 (4)

^a See text for experimental details.

25 ^b Results represent the geometric mean of 4 to 10 animals expressed in log 10 units. The number in parentheses indicates the number of animals.

^c The dose was 10 ug of each protein.

30

As demonstrated in Table 6, human RS virus fusion (and G) protein elicited very high titer of antibodies that showed high neutralizing and anti-fusion activities against BRSV. On

35

-54-

the other hand, the Diamond BRSV vaccine elected antibodies that had no detectable anti-fusion activity and significantly lower neutralization activity. Thus, the results presented in Table 6 clearly indicate that human RS virus fusion protein
5 elicits antibodies that neutralize BRSV and prevent fusion induced by BRSV. Hence, the human RS virus fusion protein elicits a protective immune response against BRSV.

6.5. AVOIDANCE OF ENHANCED DISEASE

10 Immunization or vaccination with formalin inactivated RS virus (Lot 100, Pfizer) has been associated with the development of atypical and more severe disease when immunized subjects have been subsequently infected with RS virus. (Kim et al., 1969, Am. J. Epidemiol. 89:422-34; Chin et al., 1969,
15 Am. J. Epidemiol. 89:449-63). Such potentiation of RS virus infection has been a major hurdle for the development of an efficacious and safe RS virus vaccine formulation for human and animal subjects.

Prince et al. (1985, J. Virol. 57:721-28) developed an
20 animal model for the assessment of the potentiation or enhancement of RS virus induced disease. The cotton rat animal model developed by Prince was employed to investigate whether or not similar potentiation or enhancement of subsequent disease was associated with the use of RS virus F
25 protein as an immunogen according to one embodiment of the present invention. Briefly, cotton rats were immunized by intramuscular injection of either RS virus F protein, with or without Alum as an adjuvant, or one of a variety of immunogens as indicated in Table 7. All animals received three
30 injections either on week 0, 1 and 2 or on week 0, 2, and 4, except for one group which received only one immunization. One to two weeks following the last immunization, animals were challenged with 10^4 - 10^6 PFU/animal of RS virus. At a 1 and/or

-55-

4 days post-challenge, the lungs of the animals were examined for morphological, inflammatory and pathological changes in alveolar ducts and/or alveoli according to the method of Prince et al., supra. The overall pathology of large and
5 small airways of lungs were scored using a scheme ranging from mild (+1), moderate (+2), severe (+3) to very severe (+4). Results are presented in Table 7.

10

15

20

25

30

35

-56-

Table 7

ENHANCED PATHOLOGY COTTON RAT MODEL:
FREQUENCY OF OCCLUDED SMALL AIRWAYS

5	Immunogen (Dose) ^a	Sample Challenge	Alveolar and/or Alveoli Pathology ^b				
			Day	N	+	(%)	Severity
	Lot-100 (undil)	RSV	1	2	1	(50)	1.5
	Lot-100 (undil)	RSV	4	3	2	(67)	1.5
10	RSV-formalin (undil-1:625)***	RSV	1	18	13	(72)	2.0
	RSV-formalin (undil-1:625)***	RSV	4	17	11	(65)	1.5
	RSV-live (10 ¹ -10 ⁶ PFU)	RSV	1	13	2	(15)	1.0
	RSV-live (10 ¹ -10 ⁶ PFU)	RSV	4	15	3	(20)	1.0
15	F(Alum)* (0.2-20 ug)	RSV	4	52	0	0	--
	F (5 ug)	RSV	1	3	0	0	--
	F (0.2-10 ug)	RSV	4	23	0	0	--
	F** (1-20 ug)	RSV	4	11	1	(9)	0.5
20	Medium	RSV	1	2	0		
	Medium	RSV	4	2	1	(50)	1.0
	PBS	RSV	1	4	1	(25)	1.0
	PBS	RSV	4	22	3	(14)	1.0
	None	RSV	1	3	0	0	--
25	None	RSV	4	3	0	0	--
	None*	Medium	4	3	0	0	--
	None*	None None or PBS	4	9	0	0	--

30 Footnotes continued.

-57-

a All animals were immunized by a 200 ul intramuscular injection of immunogen three times at weeks 0, 1, and 2, except for those designated * which were immunized at weeks 0, 2 and 4 and ** which received a single intramuscular injection at week 0. "****" indicates that the undiluted virus was 10^6 - 10^7 PFU.

b "Day" indicates the day post-challenge on which the pathology of lung avelolar ducts and/or alveoli were evaluated. "N" indicates the total number of animals evaluated. "+" indicates the number of animals in which pathological changes were noted, indicating enhanced or potentiated RS virus induced disease. "(%)" represents the percent of animals in which enhanced RS virus induced disease was observed. "Severity" represents the mean histopathological score on a scale of according to the method of Prince et al., supra 0 to +4)

15 As demonstrated in Table 7, animals which were immunized with RS virus fusion protein according to one embodiment of the present invention, did not exhibit pathological changes in lung tissues, including alveolar ducts and/or alveoli. On day 4, only 1 out of 86 animals immunized with fusion protein showed less than mild pathological change. On the other hand, 20 of animals immunized with commercially available RS virus vaccine Lot 100 or formalin-inactivated RS virus, greather than 65% of treated animals had mild to moderate pathological deterioration and/or inflammation of lung tissues. 25 Additionally, 20% of animals immunized with live RS virus showed at least mild pathological deterioration and/or inflamation of lung tissues. Thus, the use of RS virus fusin protein according to the present invention avoided the enhancement or potentiation of disease induced by subsequent 30 RS virus infection. Hence the vaccine formulations according to the present invention are both safe and efficacious.

-58-

7. PROTECTION OF HUMANS: RS VIRUS FUSION PROTEIN

FDA approved human phase I clinical studies were conducted to evaluate the safety and immunogenicity of RS virus fusion protein vaccine in human adult volunteers. The fusion protein from the RS virus strain A2, was purified as described in Section 6.1, supra, compounded with alum, and injected intramuscularly in the following doses: Group 1 received 5 ug (N=15), Group 2, 15 ug (N=16); and Group 3, 45 ug (N=9). One month later, 6 volunteers from Group 1 and 7 volunteers from Group 2 were boosted with a repeat injection of the respective amounts of RS virus fusion protein received at the first immunization. The schedule for vaccinations and collection of serum samples from volunteers is illustrated in Table 8.

15

Table 8DESCRIPTIVE DATA FROM PHASE I CLINICAL TRIAL OF RS VIRUS F PROTEIN IN ADULTS

Volunteers	5ug	RS Virus F Protein			Total
		15ug	45ug		
N (vax=1)	15	16	9		40
(vax=2)	6	7			13
Bleed/Vax timing:					
A(V) mean	0 days	0 days	0 days		
B mean	14.1	14.1	14.0		
C(V) mean	32.5	28.8	30.3		
D mean	61.6	55.0	58.4		
E mean	88.7	83.0	86.4		
F mean	166.0	161.4	165.1		

30

The safety data including clinical data and symptoms reported by immunized human volunteers are summarized in Table 9.

35

-59-

Table 9

PHASE-I CLINICAL TRIAL OF RS VIRUS F PROTEIN IN ADULTS:
TEMPERATURE AND LOCAL REACTIONS AT VACCINATION SITE
OCCURRING WITHIN THE FIRST 24 HOURS POST-VACCINATION

Type of Reac- tion	Score	Vaccine: Dosage:	Number of Individuals by Reaction						
			Primary				Secondary		
			5ug	15ug	45ug	All	5ug	15ug	All
10	Temper- ature	0 (<100°F)	15	15	9	39 (100%)	6	7	13 (100%)
		1 (>100°F)	0	0	0	0	0	0	0
	Redness (local)	0 (<none)	15	14	9	38 (97%)	5	7	12 (92%)
		1 (1cm)	0	1	0	1 (3%)	0	0	0
		2 (≥2cm)	0	0	0	0	1	0	1 (8%)
15	Swelling (local)	0	15	14	8	37 (94%)	6	7	13 (100%)
		1 (<1cm)	0	0	0	0	0	0	0
		2 (≥2cm)	0	0	1	2 (6%)	0	0	0
	Warmth (Local)	0	15	15	8	38 (97%)	6	7	13 (100%)
		1 (<1cm)	0	0	0	0	0	0	0
		2 (≥2cm)	0	0	1	1 (3%)	0	0	0
20	Pain (local)	0 (none)	10	6	3	19 (49%)	3	3	6 (46%)
		1 (touch/ movement)	5	6	5	16 (41%)	3	3	6 (46%)
		2 (always)	0	3	1	4 (10%)	0	1	1 (8%)

As demonstrated in Table 9, the adverse reactions (local reactions) to the immunization were very minimal and/or insignificant. Blood was drawn from the volunteers at specified time intervals and tested for the antibody titer specific to the RS virus F protein as described above. Results are presented in Table 10.

30

35

-60-

Table 10

IMMUNOGENICITY OF RS VIRUS F PROTEIN IN ADULTS EIA SEROLOGY*

Vaccination	Bleed	Geometric Mean Titer (Range) [N]		
		Group 1	Group 2	Group 3
Primary	A	66.4 (15-176) [n=15]	58.9 (21-200) [n=16]	90.9 (27-299) [n=9]
	B	160.8 ^a (30-328) [n=15]	229.0 ^a (40-965) [n=16]	415.3 ^a (179-700) [n=9]
	C	141.3 ^b (33-276) [n=15]	199.9 ^a (46-828) [n=9]	386.2 ^a (153-703) [n=9]
	D	116.2 ^c (31-217) [n=8]	235.2 ^a (73-644) [n=9]	351.1 ^b (121-645) [n=9]
	E	119.4 ^c (35-195) [n=9]	180.2 ^a (60-493) [n=9]	349.0 ^b (129-705) [n=9]
	F	106.1 (25-169) [n=9]	144.5 ^b (46-330) [n=9]	314.8 ^b (111-538) [n=9]
Boost	D	183.4 ^b (98-302) [n=6]	217.7 ^a (105-117) [n=7]	
	E	166.7 ^b (103-352) [n=6]	217.6 ^a (99-658) [n=7]	
	F	126.1 ^c (76-234) [n=6]	180.2 ^b (91-691) [n=7]	

* EIA titer is expressed as titer (X1000). Values represent geometric mean titer, with the range in parentheses and the number of samples in square brackets.

-61-

- a Comparison to Bleed A GMT by T test showed significant difference $p < 0.001$.
- b Comparison to Bleed A GMT by T test showed significant difference $p < 0.01$.
- c Comparison to Bleed A GMT by T test showed significant difference $p < 0.05$.

5

As shown in Table 10, significantly enhanced quantities of antibodies compared to the pre-existing antibodies were produced in all groups, following the immunization with either 5 or 15 or 45 ug of RS virus F protein. Hence, the RS virus F protein according to one embodiment of this invention, is very immunogenic not only in animals but also in humans.

10

8. IDENTIFICATION OF NEUTRALIZING AND/OR FUSION EPITOPE[S] OF RS VIRUS FUSION PROTEIN

15

8.1. GENERAL PROCEDURES

The following protocols were used to define an epitope of RS virus fusion protein which elicits both neutralizing and antifusion antibodies.

20

8.1.1. PROTEIN IMMUNOBLOT (WESTERN BLOT) ANALYSIS

Fusion protein subunits F_1 and F_2 were subjected to SDS-PAGE (Laemmli, 1970, Nature, 227:680-685). The separated protein subunits in the gel were electrophoretically transferred to a nitrocellulose sheet using a transfer solution containing 12.1 g Tris-HCl, 56.3 g glycine per 5 liters. The nitrocellulose sheet was air dried. The air-dried nitrocellulose was incubated at 37°C sequentially with (i) BLOTTO for 15 min., (ii) BLOTTO containing the L4 monoclonal antibody for 15 min., (iii) BLOTTO for 15 min., (iv) BLOTTO containing secondary antiserum for 60 min., and (v) BLOTTO for 15 min.

25

30

35

-62-

The secondary antiserum bound to the L4 monoclonal antibody, and was either labeled with horse radish peroxidase or not labeled. If labeled with peroxidase, antibody binding was detected by color development brought about by enzymatic reaction with 0.05% 4-chloronaphthol, 0.09% H_2O_2 in phosphate buffered saline pH 6.8. If unlabeled, antibody binding was detected by autoradiography following ^{125}I - protein A binding to the antibodies.

10 8.1.2. COUPLING OF PEPTIDES TO KEYHOLE LIMPET HEMOCYANIN (KLH) AND PRODUCTION OF RABBIT ANTISERA

A 1 mg/ml KLH solution was prepared as follows:
4 mg of a resuspended ammonium sulfate precipitate of KLH (containing 37 mg of protein/ml) was centrifuged for 5 minutes at 4°C in a microfuge. The supernatant was discarded and the pellet redissolved in 3 ml of 0.1 M $NaHCO_3$. The solution was dialyzed against 0.1 M $NaHCO_3$ with 2 changes. The volume of the dialyzed KLH solution was adjusted to 4 ml to make a 1 mg/ml KLH solution.

Four ul of a 1 mg/ml solution of a synthetic polypeptide were added to the KLH solution, and mixed at room temperature for 1 hour. Four ml of 25% aqueous glutaraldehyde were added to the mixture, and mixed for another 24 hours at room temperature. Twenty ul of 25% aqueous glutaraldehyde were again added to the mixture and mixed for a further 72 hours at room temperature. The glutaraldehyde cross-linked synthetic polypeptide - KLH was dialyzed overnight against PBS with several changes of the dialysis buffer.

Rabbits were immunized with 250 ug of protein in complete Freund's adjuvant, and boosted 2-3 times at 2 week intervals with 250 ug of protein in incomplete Freund's adjuvant.

-63-

8.1.3. PROTEOLYTIC CLEAVAGE OF FUSION PROTEINTrypsin Digestion without L4 protection:

Purified fusion protein in 50 mM Tris, pH 7.1, .05% SDS and .1% beta-mercaptoethanol was heated for 5 minutes at 100°C. Trypsin was added to the cooled, denatured fusion protein sample and incubated at 37°C for 2 hours. Enzyme to substrate ratios of 1:1000, 1:2500, and 1:5000 were used. Unless otherwise stated, all proteinases are added from a 1 mg/ml stock solution.

Trypsin Digestion with L4 protection:

Equal molar amounts of purified fusion protein and L4 monoclonal antibody were mixed and set on ice for 1 hour. Trypsin was added to the mixture in the presence of 50 mM Tris, pH 7.1, and incubated at 37°C for 2 hours. An enzyme to substrate ratio of 1:10 was used.

Trypsin/Arg-C Digestions with L4 protection:

Equal molar amounts of purified fusion protein and L4 monoclonal antibody were mixed and set on ice for 1 hour. Trypsin digestion at an enzyme to substrate ratio of 1:10 was carried out as described in Example 2.2. The trypsin-digested fusion protein was heated for 5 minutes at 100°C, cooled and digested further by Arg-C at enzyme to substrate ratios of 1:2, 1:5, 1:10 at 37°C for 2 hours.

Arg-C/Arg-C Digestions with L4 protection:

Equal molar amounts of purified fusion protein and L4 monoclonal antibody were mixed and set on ice for 1 hour. A first Arg-C digestion of the mixture at an enzyme to substrate ratio of 1:3 was performed in the presence of 20 mM NH_4HCO_3 for 2 hours at 37°C. Then the reaction mixture was reduced by adding 2-mercaptoethanol to a final concentration of 0.1% and

-64-

heated for 5 minutes at 100°C. After cooling, a second Arg-C digestion at an enzyme to substrate ratio of 1:6 for 2 hours at 37°C was carried out.

5 Lys-C/Arg-C Digestions with L4 protection:

The same protocol was followed as in Example 2.4 above except that Lys-C at an enzyme to substrate ratio of 1:2 was used in the first digestion.

10

8.1.4. DOT BLOT ANALYSIS

Dot blot analysis was performed on synthetic polypeptides as follows:

Polypeptides (up to 20 ug) were spotted on the nitrocellulose sheets. The nitrocellulose sheet was air
15 dried. The air-dried nitrocellulose was incubated at 37°C sequentially with (i) BLOTTO (5% Carnation™ instant milk in phosphate buffered saline pH 6.8) for 15 min., (ii) BLOTTO containing the L4 monoclonal antibody for 0.25-72 hrs, (iii) BLOTTO for 15 min., (iv) BLOTTO containing secondary antiserum
20 for 60-120 min., and (v) BLOTTO for 15 min.

The secondary antiserum bound to the L4 monoclonal antibody, and was either labeled with horse radish peroxidase or not labeled. If labeled with peroxidase, antibody binding was detected by color development brought about by enzymatic
25 reaction with 0.06% 4-chloronaphthol, 0.3% H₂O₂ in phosphate buffered saline pH 6.8. If unlabeled, antibody binding was detected by autoradiography following ¹²⁵I- protein A binding to the antibodies.

30

8.2. MAPPING BY DEFINED PROTEOLYTIC CLEAVAGE

In order to map the F₁ subunit of the RS virus fusion protein, synthetic polypeptides were prepared corresponding to various regions of the F₁ subunit as depicted in Figure 3 and
35

-65-

designated synthetic polypeptides (sp) sp1 through sp5. These sp were coupled to KLH as described supra and employed separately to immunize rabbits. Five corresponding rabbit antisera were obtained designated anti-sp1 through anti-sp5.

- 5 All five antisera were reactive with the F₁ subunit as well as with the particular sp used as immunogen.

The purified RS virus fusion protein was then subjected to defined proteolytic cleavage as described in detail Section 7.1.3. The cleaved protein fragments were separated by SDS-
 10 PAGE to determine molecular weight and analyzed by Western blot analysis for ability to bind to the L4 monoclonal antibody as well as to the antisera raised against the synthetic polypeptides. The position of a given proteolytic fragment within the fusion protein sequence was deduced from
 15 the reactivities with each of the antisera. Results are tabulated in Table 11 (A-E) and summarized in Figure 4.

Table 11
 MAPPING^a AND ANTIGENICITY OF PEPTIDE FRAGMENTS OF F PROTEIN

A. Trypsin Digestion of F Protein						
Fragment	I ^b	II	III	IV	V	VI
Probe	28K-1 ^c	25K-1	24K	23K	15K	14K
L4	+	+	+	+	-	+
anti-sp(1)	+	+	+	+	+	-
anti-sp(4)	+	+	+	+	+	-
anti-sp(5)	-	-	+	-	-	-
25 Predicted	137-394	137-364	137-359	156-364	137-272	
Amino Acid Position ^d						

-66-

B. Trypsin Digestion of L4-Protected F Protein

Fragment Probe	VII 28K	VIII 26K	III 24K
L4	+	-	+
5 anti-sp(1)	-	-	+
anti-sp(2)	+	+	-
anti-sp(4)	-	-	+
Predicted Amino Acid Position	272-520	336-574	137-359

C. Trypsin Digestion of L4-Protected F Protein Followed By Arg-C Digestion

Fragment Probe	III 24K	IV 23K	IX 20K	V 15K
L4	+	+	+	-
anti-sp(4)	+	+	+	+
15 Predicted Amino Acid Position	137-359	156-364	156-336	137-272

D. Arg C Digestion of F Protein

Fragment Probe	X 25K	XI 23.5K	XII 21K	XIII 14K	XIV 13K
L4	+	-	+	+	+
20 anti-sp(2)	-	+	-	-	-
anti-sp(4)	-	-	+	-	-
anti-sp(5)	-	+	-	-	-
Predicted Amino Acid Position	282-507	336-553	235-429		

E. Lys C Digestion of L4 Protected F Protein Followed By Arg C Digestion

Fragment Probe	X 25K	XV 25K	XI 23.5K	XII 21K	XVI 16K	XVII 14K	XVIII 13K
L4	+	-	-	+	-	+	+
anti-sp(2)	-	+	+	-	-	-	-
anti-sp(4)	-	-	-	+	+	-	-
30 anti-sp(5)	-	-	-	-	-	-	-
Predicted Amino Acid Position	282- 507	327- 553	156- 336	235- 429	137- 282	272- 399	282- 399

Footnotes continued.

-67-

- a The Roman Numerals identify the fragments shown in Figure 4. Approximate molecular weight (daltons) of cleavage fragments are indicated under Roman Numerals.
- b The predicted amino acid positions of the various fragments were deduced from their reactivity with the anti-synthetic polypeptide sera as well as the known amino acid specificities of the proteases.
-

Table 11 (A-E) shows the fragments which were derived by the five proteolytic cleavage protocols along with the reactivity to the monoclonal and polyclonal antisera. Figure 4 consolidates the data of Table 11 (A-E), and shows the position of all of the fragments identified in a linear map. The fragments which reacted with the L4 monoclonal antibody are shown as shaded bars and those which did not react are shown in solid bars. The cross-hatched area present in all fragments reactive with L4 monoclonal antibody spans amino acid residues 283-327.

8.3. MAPPING BY EXPRESSION OF PROTEIN FRAGMENTS

The cDNA substantially as depicted in Figure 1 containing the complete nucleotide sequence of the fusion protein gene was cloned into E. coli plasmid vector pG103 at the BamHI site. Figure 5 illustrates the resultant recombinant plasmid pPX1043.

The cDNA substantially as depicted in Figure 1 containing the complete nucleotide sequence of the fusion protein gene was cloned into E. coli plasmid vector pBR322 at the BamHI site. The resultant plasmid was designated pBR322-F. Regions of the fusion protein gene were excised from pBR322-F by restriction endonuclease digestion and ligated into the E. coli expression vector pUC19 (Yanish-Perron et al., 1985, Gene 33:103-19). Table 12 shows the restriction

-68-

sites used, the nucleotide sequences of the fusion protein gene within the restriction fragments and the amino acids encoded by these nucleotide sequences. Each of these DNA fragments was cloned into the *E. coli* expression vector pUC19 in the polylinker region such that the sequences were in frame with the lac Z gene initiation codon. The amino acid sequences encoded by the recombinant DNA molecules at and near the junctions of the vector plasmid and the cloned fusion protein gene fragments are shown in Figure 6. Since coding sequences of the fusion protein were inserted into the lac Z gene of the pUC19 vector, some amino acids from the lac Z protein as well as some amino acids encoded by the polylinker are included in the recombinant proteins.

15

Table 12
FRAGMENTS OF FUSION PROTEIN GENE CLONED
INTO THE pUC19 EXPRESSION VECTOR

Construction	Recombinant Protein	Restriction Sites Used ^a	Nucleotide Boundaries ^b	Amino Acid Boundaries
CF3	F3	HpaI-BamHI	769-1737	253-574
CF4	F4	PstI-NsiI	646-1479	212-490
CF7	F7	PstI-TaqI	646-1102	212-363
CF8	F8	PstI-HincII	646-1165	212-384
CF10	F10	PstI-NdeI	646-907	212-298
CF11	F11	NdeI-NscI	908-1479	299-490

^a A clone of the RSV fusion protein gene was cleaved with the indicated combinations of restriction endonucleases to generate RSV fragments for subsequent cloning into the expression vector pUC19. The nucleotides included within these restriction fragments are indicated along with the predicted amino acids which were expressed.

^b Nucleotide 1900 is the beginning of the synthetic BamHI linker. Nucleotides 1742-44 is a translation termination signal (TAA) so that amino acid 574 is a terminal amino acid.

35

-69-

Six recombinant proteins, designated F3, F4, F7, F8, F10, and F11, corresponding to proteins encoded by construction cF3 (also designated as plasmid pPX-1029), cF4, cF7, cF8, cF10 and cF11, were expressed in E. coli as demonstrated by reactivity to polyclonal rabbit antiserum to the native fusion proteins by Western blot analysis as described supra. All of the recombinant proteins except F11 reacted with the L4 monoclonal antibody by Western blot analysis. Figure 7 shows a diagrammatic scheme of the L4 reactive and L4 non-reactive recombinant proteins. As illustrated in Figure 7, the RS virus fusion protein sequence common to all of the recombinant proteins which are reactive with the L4 monoclonal antibody is defined by residues 253 to 298.

8.4. MAPPING BY SYNTHETIC PEPTIDES

Four synthetic polypeptides were prepared which corresponded to the amino acid residues 299-315, 294-315, 289-315, and 283-315 of the RS virus fusion protein respectively. The exact sequences of these four synthetic polypeptides are shown in Table 13.

Table 13

SEQUENCE OF SYNTHETIC POLYPEPTIDES TESTED FOR REACTIVITY WITH THE L4 MONOCLONAL ANTIBODY

Peptide	Sequence ^a	Reactivity with L4
299-315	YVVQLPLYGVIDTPCWK	-
294-315	EEVLAYVVQLPLYGVIDTPCWK	+
289-315	MSIIKEEVLAYVVQLPLYGVIDTPCWK	+
283-315	QQSYSIMSIIEVLAYVVQLPLYGVIDTPCWK	+

^a Sequences are given by the standard single letter amino acid code.

-70-

The polypeptides were spotted on nitrocellulose sheets and tested for reactivity with the L4 monoclonal antibody by dot blot analysis (see Section 7, supra).

5 Results obtained are illustrated in Figure 8. As demonstrated in Figure 8, peptide 299-315 did not react with the L4 monoclonal antibody whereas polypeptides 294-315, 289-315 and 283-315 did, indicating that a neutralizing and fusion epitope resides between residues 283-298 and can be as small
10 as 294-299. The reasons why the minimum region is set at 294-299 rather than 294-298 is because generally a stretch of at least six amino acids is necessary to form an epitope.

15 9. IMMUNOGENICITY OF MODIFIED RS VIRUS FUSION PROTEIN

9.1. CONFORMATIONAL MODIFICATION

Crosslinking experiments demonstrated that the RS virus fusion protein exists as a dimer (140,000 daltons) on the surface of virus infected cells and in substantially pure
20 preparations. In order to investigate the potential effects of protein conformation on the vaccine utility, the quaternary structure of the F protein was modified and resultant effects on immunogenicity evaluated.

A monomeric F protein (70,000 daltons) was obtained from
25 substantially pure F protein by treating the F protein at 100°C for two minutes or by treatment with 99.9% acetonitrile /0.1% TFA. The immunogenicity of dimeric and monomeric forms was tested in cotton rats. Experimental details and results are presented in Table 14.

30

35

-71-

Table 14

IMMUNOGENICITY OF RSV-F PROTEIN:
EFFECT OF PROTEIN CONFORMATION^a

Immunogen *	Dose & Route ug(times)	N	EIA log 10	Neut. Titer log 10
F-dimer	5(3x IM)	15	5.7	3.8
F-monomer	5(3x SC)	5	<2.0	<1.7

- 10 a The monomeric form was derived from native dimeric form of the F protein by heat treatment. After the treatments, cotton rats were immunized (intramuscular or subcutaneous) 3 times with the indicated immunogens at 2 week intervals and at week 6. Serum was collected and assayed for their titers.

15 The total neutralizing antibody titers were found to be 100-1000 fold less in the group that received the monomeric form, prepared by heat treatment.

20 9.2. DEACYLATED FUSION PROTEIN

In order to determine the effect of covalently attached fatty acid on the immunogenicity of RS virus, the fusion protein was deacylated using hydroxylamine (1 M for 4 hours at room temperature) and the efficacy of the deacylated protein was compared to that of F protein. Experimental details and results are presented in Table 15.

30

35

-72-

Table 15

5 IMMUNOGENICITY AND PROTECTIVE EFFICACY OF F
 PROTEIN: EFFECT OF FATTY ACID DEACYLATION^a

	Immunogen Type	Amount (ug)	N	Pre-challenge GMT		RSV isolation and titers from lungs ^b		
				EIA-F	Neut(A2)	(+)	(-)	GMT
10	F	0.5	5	156,000	233	5	0	3.36
		5.0	3	163,000	429	2	1	2.04
		deacylated						
	F	0.5	6	170,000	231	5	1	3.31
15		5.0	6	363,000	510	6	0	3.20
	PBS	0.0	6	<100	<20	6	0	4.32

20 ^a Cotton rats were immunized with the indicated immunogens at week 0, 2 and 4. At week 6, they were challenged with RSV/A2 virus intranasally. Serum was collected from the day of challenge and assayed.

25 ^b "+" represents the number of animals with detectable RS virus in their lungs at 4 day post-challenge.
 "-" represents the number of animals without detectable RS virus in their lungs at 4 days post-challenge. GMT titres are expressed as PFU/GM tissue.

30 Experimental results indicate that the protective efficacy of the deacylated F protein at 5ug dose was reduced compared to the acylated F protein.

-73-

10. PROTECTION OF ANIMALS: RS VIRUS G PROTEIN10.1. ISOLATION OF G PROTEIN

Substantially pure RS virus G protein suitable for use
5 as an immunogen in a subunit vaccine formulation was prepared
essentially according to the method of Walsh et al., 1984, J.
Gen. Virol. 65:761-67.

10.2. PASSIVE PROTECTION

10 Rabbits were immunized using the affinity purified G
protein with an intramuscular injection of 2.4 ug of RS virus
G protein in Freund's Complete Adjuvant at week 0 and with 4.8
ug of G protein in Freund's Incomplete Adjuvant at weeks 4 and
8. Animals were exsanguinated on week 17. The IgG fractions
15 of sera from the G-immunized rabbits and from a control group
of non-immunized rabbits were obtained using a protein A
Sepharese™ column.

Twenty cotton rats were divided into 3 groups of 4
animals each and two groups of 5 and 3 animals each. The
20 animals were passively immunized by intraperitoneal injection
as follows: Group 1 received 4.12 mg of rabbit anti-G IgG;
Group 2, 1.03 mg of rabbit anti-G IgG; Group 3, 4.28 mg of
normal rabbit IgG; Group 4, 1.07 mg of normal rabbit IgG; and
Group 5 (control), an equivalent volume of PBS. Twenty-four
25 hours post-immunization, animals were challenged with 5.5
log₁₀ PFU RS virus Long strain. Lungs were harvested on day 4
post-challenge. The presence and/or quantity of RS virus in
lung tissues was determined as described above. Serum samples
were collected and assays were conducted as described above.
30 Results are presented in Table 16.

-74-

Table 16

**PROTECTIVE EFFICACY OF PASSIVE
IMMUNIZATION OF ANTI-G ANTIBODIES**

Amount of IgG Administered ^a	Serology ^b		Virus Assay ^c		
	EIA	Virus Neutral- ization	+	-	TITER
Anti G-IgG 4.12 mg	5.5	3.3	0	4	<1.0
10 Anti G-IgG 1.03 mg	4.8	3.0	0	4	<1.0
Normal IgG 4.28 mg	1.0	2.0	5	0	4.5
Normal IgG 1.07 mg	1.0	2.0	3	0	4.9
none (PBS)	1.0	2.0	4	0	5.0

15 ^a Cotton rats were passively immunized using an intraperitoneal injection of rabbit immunoglobulin elicited using purified G protein from RS virus Long strain. The number of animals in each group was four, except those which received 4.28 mg normal rabbit IgG (5 animals) and 1.07 mg normal rabbit IgG (3 animals).

20 ^b Assays were performed on serum samples obtained on the day of challenge (see text for details). Results represent the geometric mean titer for 3-5 animals expressed in log₁₀ units (EIA) and log₁₀ units/ml serum (Virus Neutralization Assay).

25 ^c Long strain RS virus 5.5 log₁₀ PFU was inoculated intranasally (IN) into the passively immunized animals at 24 hours post-immunization and lungs were harvested at 4 days post-challenge. Virus was isolated from the lung tissues. "+" indicates the number of animals in each group in which virus was detectable in lungs. "--" indicates the member in which no virus was detected. "Titer" represents the geometric mean titer (PFU/gm of tissue) of virus isolated from lung tissue.

30

As demonstrated in Table 16, passive administration of rabbit antibody induced by purified G protein from RS virus

35

-75-

was efficacious in protecting cotton rats against infection by homologous RS virus.

10.3. ACTIVE IMMUNIZATION AND PROTECTION

5 Thirty seven cotton rats were divided into 3 groups of
12-13 animals each. One group was then further subdivided
into 4 groups of 3 animals each, thus forming 6 groups. All
animals in all groups, except Group 6, were immunized by
intramuscular injection of 10 ug of RS virus G protein,
10 obtained as described in Section above, in different adjuvants
as follows: Group 1 received three injections at weeks 0, 2
and 4-5 of G protein in PBS; Group 2, three injections at
weeks 0, 2 and 5 of G protein in Alum; Group 3, two injections
at weeks 0 and 5 of G protein in Alum; Group 4, three
15 injections at weeks 0, 2 and 5 of G protein in ISCOM; and
Group 5, two injections at weeks 0 and 5 of G protein in
ISCOM. Group 6, the control group, received three injections
at weeks 0, 2 and 4 of an equivalent volume of PBS alone.
Between weeks 6 and 7, all animals were challenged
20 intranasally with 4.0-5.0 log₁₀ PFU RS virus Long strain.
Serum samples were collected on the day of challenge and
assayed as described above in Section 6.2, supra. The
presence and/or quantity of RS virus in lung tissues was
determined at 4 days post-challenge as described above.
25 Results are presented in Table 17.

30

35

-76-

Table 17

IMMUNOGENICITY AND PROTECTIVE EFFICACY OF RS VIRUS G PROTEIN

Group No.	Immunogen (Adjuvant) ^a	N	EIA	Neutra- lization	Serology ^b Virus Assay ^c		
					N	+	- GMT
1	(PBS)	12	3.4	4.0	6	1	5 1.4
2	(Alum)	3	6.1	5.9	3	0	3 <1.0
3	(Alum)	3	6.3	5.3	3	0	3 <1.0
4	(ISCOM)	3	6.4	5.7	3	0	3 <1.0
5	(ISCOM)	3	6.3	5.3	3	0	3 <1.0
6	PBS	13	<1.0	<2.0	13	13	0 4.3

^a In all cases, the immunogen was 10 ug G protein in an adjuvant as indicated.

^b Assays were performed on serum samples obtained on the day of challenge. Results represent the geometric mean titer expressed in log₁₀ units.

^c Virus was isolated from the lung tissues. "N" indicates the number of animals examined. "+" indicates the number of animals in each group in which virus was detectable. "-" indicates the member in which no virus was detected. "Titer" represents the geometric mean titer (PFU/gm of tissue) of virus isolated from lung tissue.

As illustrated in Table 17, active immunization of cotton rats with G protein obtained from RS virus Long strain induced a significant immune response as demonstrated by high titers of anti-G antibody (EIA assay) that were capable of neutralizing RS virus Long strain (Neutralization assay).

Most importantly, when immunized animals were challenged with RV virus Long strain, they were effectively protected against such infection as compared to control animals.

-77-

11. IMMUNOGENICITY OF RS VIRUS G PROTEIN
EXPRESSED IN RECOMBINANT VECTORS

A recombinant vector expressing the RS virus Strain A2 G protein was prepared. Plasmid pPL-lambda (pPL), which carries the P_L promoter and the N gene on a 1215 bp segment of the bacteriophage lambda genome was purchased from Pharmacia Fine Chemicals (Piscataway, NJ). A unique HpaI site in the N gene is located 321 bp downstream from the start of P_L transcription. Sequences inserted into this restriction site will thus be regulated by the P_L promoter. An expression vector plasmid pPX1600 was constructed by inserting a synthetic oligonucleotide sequence into the HpaI site of the pPL plasmid. This oligonucleotide contained a translation termination codon in frame with the N coding sequences followed by translation initiation signals (Shine-Dalgarno box plus ATG) and unique restriction sites (NcoI, StuI, EcoRV) for insertion of heterologous DNA sequences in frame with the synthetic ATG. A full length cDNA corresponding to the G glycoprotein gene that was flanked by BamHI linkers (Elango et al., 1980, Proc. Nat'l Acad. Sci. USA 83:1906-10) was cloned into the BamHI site of pBR322. After BamHI digestion, the RS virus gene encoding the G protein was excised from pBR322, filled with the Klenow fragments of DNA polymerase I, and blunt end ligated to StuI-cleaved pPX1600. Following transformation of E. coli (N99cI⁺ strain), the plasmid pPX1044 was isolated (Figure 9). In this plasmid, the RS virus G encoding sequence was in the correct orientation and reading frame for P_L -directed expression by readthrough from the synthetic ATG.

Plasmid pPX1044 was introduced into Salmonella typhimurium (LB50150) cells by the $CaCl_2$ procedure. The transformed cells were grown to late log phase and the expressed RS virus G protein was purified by an immunoaffinity

-78-

method using L9 monoclonal antibody (Walsh and Hruska, 1983 J. Virol. 47:171-177). The recombinant non-glycosylated G protein produced significant neutralization titers (3.7 logs/ml against Long strain) in rabbits, following 4 immunizations with 10 ug in the presence of Freund's adjuvant. A similar neutralizing antibody response was elicited in cotton rats after 3 immunizations with 5ug of the recombinant non-glycosylated G protein. Hence, the bacterially derived recombinant G is immunogenic and the antibodies produced against the recombinant G are functional and therefore will be protective against RS virus infection.

12. CELL-MEDIATED IMMUNOLOGICAL ASPECTS OF RS VIRUS VACCINE

The data presented in this application demonstrate that immunization with RSV glycoprotein(s) either F protein alone or in combination with G protein, induces circulating antibodies that are functional and protective. In addition to this antibody response, cell mediated immunity may be of importance for the protective efficacy. Indeed, several studies have shown that many viral infections are associated with the induction of cytotoxic T cells which recognize viral glycoproteins. Cytotoxic T cells against RS virus have been described in mouse and humans (Taylor, et al., 1984, Infect. Immun. 43:649-55; Bangham et al., 1985, J. Virol 56:55-59; Bangham et al., 1986, Proc. Nat'l Acad Sci. USA 83:9183-87). Moreover, it has been shown that passive transfer of T cells from RSV-primed mice can clear a presistent RSV infection when the cells are administered to athymic nude mice (Cannon et al., 1987, Immunol. 62:133-38). In order to investigate whether immunization with RS virus F protein could induce cytotoxic T cell responses, mice were immunized with purified RS virus F protein and immune-stimulated effector T cells were assayed in vitro. Responses in animals immunized with live RS

-79-

virus were also examined. In practice, 36 mice were divided into 4 groups and animals were immunized at 0 and 2 months as follows: Group 1 received 2×10^6 TCID50 Long strain RS virus intranasal and intraperitoneal injections respectively; Group 2, 20 ug RS virus F protein (in Alum) via intranasal injections; Group 3, 20 ug RS virus F protein (in Alum) via intramuscular injections; and Group 4, were a non-immunized control group. Three week after the last booster injection, animals were sacrificed and spleens were harvested. Effector T cells were obtained from the spleens and were re-stimulated in vitro with either peritoneal exudate cells (PEC) or spleen cells infected with RS virus. Cytotoxic T cells were tested using a Cr-51 release assay (Bangham et al., 1985, J. Virol. 56:55-59). The results are presented in Table 18.

15

Table 18

CTL RESPONSE INDUCED BY RSV-F PROTEIN

20	Group No. ^a	In Vitro Stimulation	% Cr Release From Target Cells ^b					
			RSV + PEC		BCH-4		RSV - PEC ^c	
			40:1	10:1	40:1	10:1	40:1	10:1
25	1	RSV+PEC	10.1	3.9	6.2	-4.2	0.8	0.2
		RSV+spleen	23.8	10.0	15.2	-2.0	-0.2	1.1
		None	-0.2	--	-0.2	--	-0.3	--
	2	RSV+PEC	12.4	4.0	9.2	-2.6	0.97	-0.8
		RSV+spleen	-0.2	-0.2	-1.4	-0.6	-0.1	1.5
		None	-0.7	--	1.6	--	-0.8	--
	3	RSV+PEC	2.4	2.4	8.4	1.2	0.4	-0.8
		RSV+spleen	-0.3	-0.1	-6.0	-5.3	0.8	-1.6
		None	-0.2	--	0.2	--	-5.8	--
30	4	RSV+PEC	0.4	--	5.8	--	0.5	--
		RSV+spleen	-0.4	--	1.0	--	-0.2	--
		None	-1.5	--	0.6	--	0.2	--

Footnotes continued.

35

-80-

- a See text for experimental details.
- b Target cells were BCH₄ cells (a persistently RSV Long strain infected Balb/c fibroblast cell line), RSV infected peritoneal exudate cells (PEC) and uninfected PEC cells.

- 5 c The ratio indicates the ratio of effector: target cells.
-

10 Significant levels of RS virus-specific cytotoxic T cells were induced following immunization with RS virus F protein (IP/IP) i.e. Group 2 and with RS virus (IN/IP), i.e., Group 1. No significant T cell response was observed in animals which received RS virus F protein via an intramuscular route, i.e., Group 3 or in control animals, i.e., Group 4. The induction of cytotoxic T cells thus depends upon the route of
15 inoculation, proteins versus live virus and the cells used for restimulation. Purified F protein of RSV is able to induce not only humoral but also cellular immunity.

20 13. DEPOSIT OF MICROORGANISMS

Many polynucleotide sequences may be used to practice the present invention. E. coli strain JM83 carrying plasmid pPX1043 which comprises the complete RS Virus fusion protein gene and an E. coli strain JM103 carrying plasmid pPX1029 have been deposited with the Agricultural Research Culture
25 Collection (NRRL), Peoria, IL and have been assigned accession numbers NRRL B-18254 and NRRL B-18255 respectively. E. coli strain N99cI⁺ carrying plasmid pPX1044 has also been deposited with the Agricultural Research Culture Collection (NRRL),
30 Peoria, IL and has been assigned accession number B-18419.

-81-

The present invention is not to be limited in scope by the microorganisms deposited; since the deposit of microorganisms is only intended as a single illustration of one aspect of the invention.

5 Many variations of this invention as herein set forth may be made without departing from the spirit and scope thereof. Such variations are intended to fall within the scope of the appended claims. The specific embodiments described are given by way of example only, and the invention
10 is limited only by the appended claims.

15

20

25

30

35

-83-

PCT Form 134 continued

<u>Deposited Microorganism</u>	<u>Accession Number</u>	<u>Date of Deposit</u>
<u>Eschirichia coli</u> JM 83 carrying pPx1043	B-18254	September 30, 1987
<u>Eschirichia coli</u> JM 103 carrying pPx1029	B-18255	September 30, 1987

Depository Institution: Agricultural Research Culture Collection
1815 North University Street
Peoria, Illinois 61604

-84-

WHAT IS CLAIMED IS:

1. A substantially pure polypeptide related to a neutralizing and/or fusion epitope of Respiratory Syncytial virus fusion protein.
5
2. The polypeptide according to claim 1 having a molecular weight of about 140,000 daltons.
- 10 3. The polypeptide according to claim 1 having an amino acid sequence comprising the sequence substantially as depicted in Figure 1 from amino acid residue 137 to residue 574 or any antigenic portion thereof.
- 15 4. The polypeptide according to claim 1 having an amino acid sequence comprising the sequence substantially as depicted in Figure 1 from amino acid residue 253 to residue 298.
- 20 5. The polypeptide according to claim 1 having an amino acid sequence comprising the sequence substantially as depicted in Figure 1 from amino acid residue 283 to residue 315.
- 25 6. The polypeptide according to claim 1 having an amino acid sequence comprising the sequence substantially as depicted in Figure 1 from amino acid residue 289 to residue 315.
- 30 7. The polypeptide according to claim 1 having an amino acid sequence comprising the sequence substantially as depicted in Figure 1 from amino acid residue 294 to residue 315.

-85-

8. The polypeptide according to claim 1 having a molecular weight of about 700 to about 4000 daltons which polypeptide binds to a monoclonal antibody which has Respiratory Syncytial Virus neutralizing activity.

5

9. The polypeptide according to claim 1 having a molecular weight between about 700 to about 4000 daltons which polypeptide binds to a monoclonal antibody which has Respiratory Syncytial Virus neutralizing activity and anti-
10 Respiratory Syncytial Virus induced fusion activity.

10. The polypeptide according to claim 1 having a molecular weight between about 700 to about 4000 daltons which polypeptide binds to a monoclonal antibody which has anti-
15 Respiratory Syncytial Virus induced fusion activity.

11. The polypeptide according to claim 1 having palmitate covalently attached thereto.

20 12. The polypeptide according to claim 2 having palmitate covalently attached thereto.

13. A substantially pure polypeptide, comprising a polypeptide having a molecular weight af about 700 to about
25 4000 daltons and having the amino acid sequence Gln - Gln - Ser - Tyr - Ser - Ile - Met - Ser - Ile - Ile - Lys - Glu - Glu - Val - Leu - Ala - Tyr - Val - Val - Gln - Leu - Pro - Leu - Tyr - Gly - Val - Ile - Asp - Thr - Pro - Cys - Trp - Lys.

30

35

-86-

14. A substantially pure polypeptide, comprising a polypeptide having a molecular weight between about 700 to about 4000 daltons and having the amino acid sequence Met - Ser - Ile - Ile - Lys - Glu - Glu - Val - Leu - Ala - Tyr - Val - Val - Gln - Leu - Pro - Leu - Tyr - Gly - Val - Ile - Asp - Thr - Pro - Cys - Trp - Lys.
15. A substantially pure polypeptide, comprising a polypeptide having a molecular weight between about 700 to about 2500 daltons and having the amino acid sequence Glu - Glu - Val - Leu - Ala - Tyr - Val.
16. A substantially pure unglycosylated polypeptide related to a neutralizing epitope of Respiratory Syncytial Virus G protein.
17. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 1, 2 or 3 mixed with a suitable pharmaceutical carrier.
18. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 4, 5 or 6 mixed with a suitable pharmaceutical carrier.
19. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 7, 8 or 9 mixed with a suitable pharmaceutical carrier.
20. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 10, 11 or 12 mixed with a suitable pharmaceutical carrier.

-87-

21. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 13, 14, 15 or 16 mixed with a suitable pharmaceutical carrier.

5

22. A subunit vaccine formulation, in which the immunogen comprises an effective amount of the polypeptide of claim 1 mixed with a substantially pure polypeptide related to a neutralizing epitope of Respiratory Syncytial Virus G protein and a suitable pharmaceutical carrier.

10

23. A subunit vaccine formulation in which the immunogen comprises an effective amount of a substantially pure non-glycosylated polypeptide related to a neutralizing epitope of Respiratory Syncytial Virus G protein.

15

24. The subunit vaccine formulation according to claim 23, further comprising a pharmaceutical carrier.

25. A method of actively immunizing an animal or a human against Respiratory Syncytial Virus, comprising administering an effective amount of the polypeptide of claim 1.

20

26. A method of actively immunizing an animal or a human against Respiratory Syncytial Virus, comprising administering an effective amount of the polypeptide of claim 1 in combination with an effective amount of a substantially pure polypeptide related to a neutralizing epitope of Respiratory Syncytial Virus G protein and a suitable pharmaceutical carrier.

25

30

35

-88-

27. A recombinant virus the genome of which comprises a nucleotide sequence encoding a neutralizing and/or fusion epitope of Respiratory Syncytial Virus fusion protein or an antigenic portion thereof which is under control of a second
5 nucleotide sequence that regulates gene expression such that a peptide or protein related to the neutralizing and/or fusion epitope of Respiratory Syncytial Virus fusion protein is expressed in a host infected with the virus.
- 10 28. The recombinant virus according to claim 27 in which the nucleotide sequence comprises the nucleotide sequence substantially as depicted in Figure 1 from amino acid residue 137 to residue 574.
- 15 29. The recombinant virus according to claim 27 in which the nucleotide sequence comprises the nucleotide sequence substantially as depicted in Figure 1 from amino acid residue 253 to residue 298.
- 20 30. The recombinant virus according to claim 27 in which the nucleotide sequence comprises the nucleotide sequence substantially as depicted in Figure 1 from amino acid residue 283 to residue 315.
- 25 31. The recombinant virus according to claim 27 in which the nucleotide sequence comprises the nucleotide sequence substantially as depicted in Figure 1 from amino acid residue 289 to residue 315.
- 30 32. The recombinant virus according to claim 27 in which the nucleotide sequence comprises the nucleotide sequence substantially as depicted in Figure 1 from amino acid residue 294 to residue 315.

-89-

33. A live virus vaccine formulation comprising a virus according to claim 27, 28, 29, 30, 31 or 32, in which the residue is infectious without causing disease in a host to be vaccinated.

5

34. An inactivated virus vaccine formulation, comprising an effective amount of the virus according to claim 27, 28, 29, 30, 31 or 32 in a non-infectious state mixed with a suitable pharmaceutical carrier.

10

35. A recombinant DNA vector, comprising substantially pPX1029.

36. A recombinant DNA vector, comprising substantially
15 pPX1044.

37. A unicellular organism containing the recombinant DNA vector of claim 35.

20 38. A unicellular organism containing the recombinant DNA vector of claim 36.

39. A bacterium containing the recombinant DNA vector of claim 35.

25

40. A bacterium containing the recombinant DNA vector of claim 36.

30 41. The bacterium of claim 37, comprising Escherichia coli deposited with the NRRL and assigned accession number B-18255 or a mutant, recombinant or genetically engineered derivative thereof.

35

-90-

42. A vaccine formulation, comprising a recombinant attenuated enteroinvasive bacterium containing a DNA sequence which encodes a neutralizing and/or fusion epitope of Respiratory Syncytial Virus fusion protein, which sequence is expressed by the bacterium, and in which the bacterium is infectious without causing significant disease in a host to be vaccinated.

43. A vaccine formulation, comprising a recombinant attenuated enteroinvasive bacterium containing a DNA sequence which encodes a neutralizing epitope of Respiratory Syncytial Virus G protein, which sequence is expressed by the bacterium, and in which the bacterium is infectious without causing significant disease in a host to be vaccinated.

44. A method for the detection of Respiratory Syncytial Virus in a sample of tissue or body fluids, comprising: using the polypeptide according to claim 1, 2, 3, 4, 5, 6, 7 or 16 as an antigen in an assay system selected from the group consisting of an enzyme-linked immunosorbent assay, a radioimmunoassay, a gel diffusion precipitation reaction assay, an immunodiffusion assay, an agglutination assay, a fluorescent immunoassay, a protein A immunoassay and an immunoelectrophoresis assay; and detecting any reaction with the antigen, in which any reaction indicates the presence of Respiratory Syncytial Virus in the sample.

30

35

-91-

45. A method for the detection of Respiratory Syncytial virus in a sample of tissue or body fluids, comprising: using a nucleotide sequence of a gene encoding a polypeptide of protein related to a fusion protein of Respiratory Syncytial virus as depicted in Figure 1 or a fragment thereof as a probe in a nucleic acid hybridization assay and detecting any reaction with the probe, in which any reaction indicates the presence of Respiratory Syncytial Virus.

46. The method according to claim 40, in which the nucleic acid hybridization assay is selected from the group consisting of Southern blots, Northern blots and colony blots.

47. The method according to claim 41, in which the nucleic acid hybridization assay is selected from the group consisting of Southern blots, Northern blots and colony blots.

20

25

30

35

1/11

FIG. 1

G GGG CAA ATA ACA ATG GAG TTG															22
Met Glu Leu															3
CTA ATC CTC AAA GCA AAT GCA AAT ACC ACA ATC CTC ACT GCA GTC	67														
Leu Ile Leu Lys Ala Asn Ala Ile Thr Thr Ile Leu Thr Ala Val	18														
ACA TTT TGT TTT GCT TCT GGT CAA AAC ATC ACT GAA GAA TTT TAT	112														
Thr Phe Cys Phe Ala Ser Gly Gln Asn Ile Thr Glu Glu Phe Tyr	33														
CAA TCA ACA TGC AGT GCA GTT AGC AAA GGC TAT CTT AGT GCT CTG	157														
Gln Ser Thr Cys Ser Ala Val Ser Lys Gly Tyr Leu Ser Ala Leu	48														
AGA ACT GGT TGG TAT ACC AGT GTT ATA ACT ATA GAA TTA AGT AAT	202														
Arg Thr Gly Trp Tyr Thr Ser Val Ile Thr Ile Glu Leu Ser Asn	63														
ATC AAG GAA AAT AAG TGT AAT GGA ACA GAT GCT AAG GTA AAA TTG	247														
Ile Lys Glu Asn Lys Cys Asn Gly Thr Asp Ala Lys Val Lys Leu	78														
ATA AAA CAA GAA TTA GAT AAA TAT AAA AAT GCT GTA ACA GAA TTG	292														
Ile Lys Gln Glu Leu Asp Lys Tyr Lys Asn Ala Val Thr Glu Leu	93														
CAG TTG CTC ATG CAA AGC ACA CCA CCA ACA AAC AAT CGA GCC AGA	337														
Gln Leu Leu Met Gln Ser Thr Pro Pro Thr Asn Asn Arg Ala Arg	108														
AGA GAA CTA CCA AGG TTT ATG AAT TAT ACA CTC AAC AAT GCC AAA	382														
Arg Glu Leu Pro Arg Phe Met Asn Tyr Thr Leu Asn Asn Ala Lys	123														
AAA ACC AAT GTA ACA TTA AGC AAG AAA AGG AAA AGA AGA TTT CTT	427														
Lys Thr Asn Val Thr Leu Ser Lys Lys Arg Lys Arg Arg Phe Leu	138														
GGT TTT TTG TTA GGT GTT GGA TCT GCA ATC GCC AGT GGC GTT GCT	472														
Gly Phe Leu Leu Gly Val Gly Ser Ala Ile Ala Ser Gly Val Ala	153														
GTA TCT AAG GTC CTG CAC CTA GAA GGG GAA GTG AAC AAG ATC AAA	517														
Val Ser Lys Val Leu His Leu Glu Gly Glu Val Asn Lys Ile Lys	168														
AGT GCT CTA CTA TCC ACA AAC AAG GCT GTA GTC AGC TTA TCA AAT	562														
Ser Ala Leu Leu Ser Thr Asn Lys Ala Val Val Ser Leu Ser Asn	183														
GGA GTT AGT GTC TTA ACC AGC AAA GTG TTA GAC CTC AAA AAC TAT	607														
Gly Val Ser Val Leu Thr Ser Lys Val Leu Asp Leu Lys Asn Tyr	198														
ATA GAT AAA CAA TTG TTA CCT ATT GTG AAC AAG CAA AGC TGC AGC	652														
Ile Asp Lys Gln Leu Leu Pro Ile Val Asn Lys Gln Ser Cys Ser	213														
ATA TCA AAT ATA GAA ACT GTG ATA GAG TTC CAA CAA AAG AAC AAC	697														
Ile Ser Asn Ile Glu Thr Val Ile Glu Phe Gln Gln Lys Asn Asn	228														
AGA CTA CTA GAG ATT ACC AGG GAA TTT AGT GTT AAT GCA GGT GTA	742														
Arg Leu Leu Glu Ile Thr Arg Glu Phe Ser Val Asn Ala Gly Val	243														
ACT ACA CCT GTA AGC ACT TAC ATG TAA ACT AAT AGT GAA TTA TTG	787														
Thr Thr Pro Val Ser Thr Tyr Met Leu Thr Asn Ser Glu Leu Leu	258														

SUBSTITUTE SHEET

2/11

FIG. 1 con't

TCA	TTA	ATC	AAT	GAT	ATG	CCT	ATA	ACA	AAT	GAT	CAG	AAA	AAG	TTA	832
Ser	Leu	Ile	Asn	Asp	Met	Pro	Ile	Thr	Asn	Asp	Gln	Lys	Lys	Leu	273
ATG	TCC	AAC	AAT	GTT	CAA	ATA	GTT	AGA	CAG	CAA	AGT	TAC	TCT	ATC	877
Met	Ser	Asn	Asn	Val	Gln	Ile	Val	Arg	Gln	Gln	Ser	Tyr	Ser	Ile	288
ATG	TCC	ATA	ATA	AAA	GAG	GAA	GTC	TTA	GCA	TAT	GTA	GTA	CAA	TTA	922
Met	Ser	Ile	Ile	Lys	Glu	Glu	Val	Leu	Ala	Tyr	Val	Val	Gln	Leu	303
CCA	CTA	TAT	GGT	GTT	ATA	GAT	ACA	CCC	TGT	TGG	AAA	CTA	CAC	ACA	967
Pro	Leu	Tyr	Gly	Val	Ile	Asp	Thr	Pro	Cys	Trp	Lys	Leu	His	Thr	318
TCC	CCT	CTA	TGT	ACA	ACC	AAC	ACA	AAA	GAA	GGG	TCC	AAC	ATC	TGT	1012
Ser	Pro	Leu	Cys	Thr	Thr	Asn	Thr	Lys	Glu	Gly	Ser	Asn	Ile	Cys	333
TTA	ACA	AGA	ACT	GAC	AGA	GGA	TGG	TAC	TGT	GAC	AAT	GCA	GGA	TCA	1057
Leu	Thr	Arg	Thr	Asp	Arg	Gly	Trp	Tyr	Cys	Asp	Asn	Ala	Gly	Ser	348
GTA	TCT	TTC	TTC	CCA	CAA	GCT	GAA	ACA	TGT	AAA	GTT	CAA	TCA	AAT	1102
Val	Ser	Phe	Phe	Pro	Gln	Ala	Glu	Thr	Cys	Lys	Val	Gln	Ser	Asn	363
CGA	GTA	TTT	TGT	GAC	ACA	ATG	AAC	AGT	TTA	ACA	TTA	CCA	AGT	GAA	1147
Arg	Val	Phe	Cys	Asp	Thr	Met	Asn	Ser	Leu	Thr	Leu	Pro	Ser	Glu	378
ATA	AAT	CTC	TGC	AAT	GTT	GAC	ATA	TTC	AAC	CCC	AAA	TAT	GAT	TGT	1192
Ile	Asn	Leu	Cys	Asn	Val	Asp	Ile	Phe	Asn	Pro	Lys	Tyr	Asp	Cys	393
AAA	ATT	ATG	ACT	TCA	AAA	ACA	GAT	GTA	AGC	AGC	TCC	GTT	ATC	ACA	1237
Lys	Ile	Met	Thr	Ser	Lys	Thr	Asp	Val	Ser	Ser	Ser	Val	Ile	Thr	408
TCT	CTA	GGA	GCC	ATT	GTG	TCA	TGC	TAT	GGC	AAA	ACT	AAA	TGT	ACA	1282
Ser	Leu	Gly	Ala	Ile	Val	Ser	Cys	Tyr	Gly	Lys	Thr	Lys	Cys	Thr	423
GCA	TCC	AAT	AAA	AAT	CGT	GGA	ATC	ATA	AAG	ACA	TTT	TCT	AAC	GGG	1327
Ala	Ser	Asn	Lys	Asn	Arg	Gly	Ile	Ile	Lys	Thr	Phe	Ser	Asn	Gly	438
TGC	GAT	TAT	GTA	TCA	AAT	AAA	GGG	ATG	GAC	ACT	GTG	TCT	GTA	GGT	1372
Cys	Asp	Tyr	Val	Ser	Asn	Lys	Gly	Met	Asp	Thr	Val	Ser	Val	Gly	453
AAC	ACA	TTA	TAT	TAT	GTA	AAT	AAG	CAA	GAA	GGT	AAA	AGT	CTC	TAT	1417
Asn	Thr	Leu	Tyr	Tyr	Val	Asn	Lys	Gln	Glu	Gly	Lys	Ser	Leu	Tyr	468
GTA	AAA	GGT	GAA	CCA	ATA	ATA	AAT	TTC	TAT	GAC	CCA	TTA	GTA	TTC	1462
Val	Lys	Gly	Glu	Pro	Ile	Ile	Asn	Phe	Tyr	Asp	Pro	Leu	Val	Phe	483
CCC	TCT	GAT	GAA	TTT	GAT	GCA	TCA	ATA	TCT	CAA	GTC	AAC	GAG	AAG	1507
Pro	Ser	Asp	Glu	Phe	Asp	Ala	Ser	Ile	Ser	Gln	Val	Asn	Glu	Lys	498
ATT	AAC	CAG	AGC	CTA	GCA	TTT	ATT	CGT	AAA	TCC	GAT	GAA	TTA	TTA	1552
Ile	Asn	Gln	Ser	Leu	Ala	Phe	Ile	Arg	Lys	Ser	Asp	Glu	Leu	Leu	513

SUBSTITUTE SHEET

3/11

FIG. 1 con't

CAT AAT GTA AAT GCT GGT AAA TCC ACC ACA AAT ATC ATG ATA ACT 1597
His Asn Val Asn Ala Gly Lys Ser Thr Thr Asn Ile Met Ile Thr 528

ACT ATA ATT ATA GTG ATT ATA GTA ATA TTG TTA TCA TTA ATT GCT 1642
Thr Ile Ile Ile Val Ile Ile Val Ile Leu Leu Ser Leu Ile Ala 543

GTT GGA CTG CTC TTA TAC TGT AAG GCC AGA AGC ACA CCA GTC ACA 1687
Val Gly Leu Leu Leu Tyr Cys Lys Ala Arg Ser Thr Pro Val Thr 558

CTA AGC AAA GAT CAA CTG AGT GGT ATA AAT AAT ATT GCA TTT AGT 1732
Leu Ser Lys Asp Gln Leu Ser Gly Ile Asn Asn Ile Ala Phe Ser 573

AAC TAA ATA AAA ATA GCA CCT AAT CAT GTT CTT ACA ATG GTT TAC 1777
Asn ***
574

TAT CTG CTC ATA GAC AAC CCA TCT GTC ATT GGA TTT TCT TAA AAT 1822

CTG AAC TTC ATC GAA ACT CTC ATC TAT AAA CCA TCT CAC TTA CAC 1867

TAT TTA AGT AGA TTC CTA GTT TAT AGT TAT AT
1899

4/11





FIG. 2

SUBSTITUTE SHEET


5/11

FIG. 3




NAME	RESIDUES	SEQUENCE
SP1	155-169	S-K-V-L-H-L-E-G-E-V-N-K-I-K-S
SP2	500-512	N-Q-S-L-A-F-I-R-K-S-D-E-L
SP3	414-429	V-S-C-Y-G-K-T-K-C-T-A-S-N-K-N-R
SP4	261-273	I-N-D-M-P-I-T-N-D-Q-K-K-L
SP5	320-340	P-L-C-T-T-N-T-L-E-G-S-N-I-C-L-T-R-T-D-R-G

6/11



7/11

FIG. 5



SUBSTITUTE SHEET

8/11

FIG. 6

A. AMINO TERMINI


<u>CONSTRUCTION</u>	<u>LAC Z</u>	<u>POLYLINKER</u>	<u>FUSION PROTEIN</u>
F3	NH ₂ -M-T-M-I-T-	P-S-L-H-S-C-W-S-	T-N-S...
F4, F8, F7, F10	NH ₂ -M-T-M-I-T-	P-S-L-H-S-	C-S-I...
F11	NH ₂ -M-T-M-I-T-	P-S-L-H-S-	Y-V-V...

B. CARBOXY TERMINI

<u>CONSTRUCTION</u>	<u>F</u>	<u>POLYLINKER</u>	<u>LAC Z</u>
F3	...-F-S-N-COOH		
F4, F11	...-F-D-A-	G-R-L-COOH	
F8	...-C-N-V-	D-S-R-G-S-P-G-T-E-	L-E-F-T-G-R-R-K-T-T-S-COOH
F7	...-Q-S-N-	R-P-A-G-R-L-COOH	
F10	...-V-L-A-COOH		

SUBSTITUTE SHEET

FIG. 7



SUBSTITUTE SHEET

10/11




FIG. 8

SUBSTITUTE SHEET

11/11




FIG. 9

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US88/03399

I. CLASSIFICATION		SUBJECT MATTER (if several classification symbols apply, indicate all) ³	
According to International Patent Classification (IPC) or to both National Classification and IPC			
INT. CL. 4th Ed. C12Q 1/70, 1/68; C12N 15/00, 7/00, 1/20; see attn: US. CL. 435/5, 6, 7, 172.3, 235, 253, 320; 436/515, 516, 533, 534, 828; A			
II. FIELDS SEARCHED			
Minimum Documentation Searched ⁴			
Classification System ¹		Classification Symbols	
US		435/5, 6, 7, 172.3, 253, 320; 235; 436/515, 516, 533, 534, 828; 935/12, 73, 78	
Documentation Searched other than Minimum Documentation to the extent that such documents are included in the fields searched ⁵			
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴			
Category ⁶	Citation of Document, ¹⁶ with indication, where appropriate, of the relevant passages ¹⁷		Relevant to Claim No. ¹⁸
X	Biological Abstract, Vol. 82, No. 9, Issued 01 November 1986 (Murphy) "Dissociation between serum neutralizing and glycoprotein antibody responses of infants and children who received inactivated respiratory syncytial virus vaccine." Abstract no. 82914		1-47
X, P	Biological Abstract, Vol. 85, No. 1, Issued 01 January 1988 (Trudel) "Identification of a synthetic peptide as part of a major neutralization epitope of respiratory syncytial virus". Abstract no. 4479		1-47
X	Canadian Journal Microbiological, Vol. 32, Issued 1986 (Trudel) "Immunovirological studies on human respiratory syncytial virus structural proteins." pages 15-21 (See abstract).		1-47
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁵ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p> </div> </div>			
IV. CERTIFICATION			
Date of the Actual Completion of the International Search ¹		Date of Mailing of this International Search Report ²	
12 January 1989		28 FEB 1989	
International Searching Authority ¹		Signature of Authorized Officer ²⁰	
ISA/US		DELBERT R. PHILLIPS	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, 1- with indication, where appropriate, of the relevant passages 1-7	Relevant to Claim No 1*
X,P	Canadian Journal Microbiology, Vol.33, Issued 1987(Trudel)"Respiratory syncytial virus fusion glycoprotein: further characterization of a major epitope involved in virus neutralization". pages 933-938(see abstract).	1-47
X	Journal of Clinical Microbiology, Vol. 23, No. 3, Issued March 1986(Anderson)"Identification of Epitopes on Respiratory Syncytial Virus Proteins by Competitive Binding Immunoassay". pages 475-480(see abstract).	1-47
X	Journal General Virology, Vol. 66, Issued 1985(Mufson)"Two Distinct Subtypes of Human Respiratory Syncytial Virus."pages 2111-2124 (see summary).	1-47
X	Journal General Virology, Vol. 67, Issued 1986, (Samson)"Neutralizing Monoclonal Antibody to Respiratory Syncytial Virus which Binds to Both F ₁ and F ₂ Components of the Fusion Protein". pages 1479-1483, (see Summary).	1-47
X	Journal General Virology, Vol. 67, Issued 1986,(Gimenez)"Antigenic Variation between Human Respiratory Syncytial Virus Isolates." pages 863-870 (See Summary).	1-47
X	Journal General Virology, Vol. 67, Issued 1986, (Walsh)"Analysis of the Respiratory Syncytial Virus Fusion Protein Using Monoclonal and Polyclonal Antibodies." pages 505-513(see Summary).	1-47
X	Journal General Virology, Vol. 66, Issued 1985, (Walsh)"Purification and Characterization of the Respiratory Syncytial Virus Fusion Protein."pages 409-415(See Summary).	1-47

PCT/US88/03399

ATTACHMENT TO PCT/ISA/210

I. CLASSIFICATION OF SUBJECT MATTER

US. CL. 935/12,73,78

THIS PAGE BLANK (USPTO)